

ON SIMPLE MODULES
FOR CERTAIN POINTED HOPF ALGEBRAS

A Dissertation

by

MARIANA PEREIRA

Submitted to the Office of Graduate Studies of
Texas A&M University
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

December 2006

Major Subject: Mathematics

ON SIMPLE MODULES
FOR CERTAIN POINTED HOPF ALGEBRAS

A Dissertation

by

MARIANA PEREIRA

Submitted to the Office of Graduate Studies of
Texas A&M University
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

Approved by:

Chair of Committee,	Sarah Witherspoon
Committee Members,	Marcelo Aguiar
	Andreas Klappenecker
	Frank Sottile
Head of Department,	Al Boggess

December 2006

Major Subject: Mathematics

ABSTRACT

On Simple Modules

for Certain Pointed Hopf Algebras. (December 2006)

Mariana Pereira, B.S., Universidad de la República, Uruguay;

M.S., University of Massachusetts

Chair of Advisory Committee: Dr. Sarah Witherspoon

In 2003, Radford introduced a new method to construct simple modules for the Drinfel'd double of a graded Hopf algebra. Until then, simple modules for such algebras were usually constructed by taking quotients of Verma modules by maximal submodules. This new method gives a more explicit construction, in the sense that the simple modules are given as subspaces of the Hopf algebra and one can easily find spanning sets for them. I use this method to study the representations of two types of pointed Hopf algebras: restricted two-parameter quantum groups, and the Drinfel'd double of rank one pointed Hopf algebras of nilpotent type. The groups of group-like elements of these Hopf algebras are abelian; hence, they fall among those Hopf algebras classified by Andruskiewitsch and Schneider. I study, in particular, under what conditions a simple module can be factored as the tensor product of a one dimensional module with a module that is naturally a module for a special quotient. For restricted two-parameter quantum groups, given θ a primitive ℓ th root of unity, the factorization of simple $\mathfrak{u}_{\theta^y, \theta^z}(\mathfrak{sl}_n)$ -modules is possible, if and only if $\gcd((y - z)n, \ell) = 1$. I construct simple modules using the computer algebra system SINGULAR::PLURAL and present computational results and conjectures about bases and dimensions. For rank one pointed Hopf algebras, given the data $\mathcal{D} = (G, \chi, a)$, the factorization of simple $D(H_{\mathcal{D}})$ -modules is possible if and only if $|\chi(a)|$ is odd and $|\chi(a)| = |a| = |\chi|$. Under this condition, the tensor product of two simple $D(H_{\mathcal{D}})$ -

modules is completely reducible, if and only if the sum of their dimensions is less or equal than $|\chi(a)| + 1$.

To my family

ACKNOWLEDGMENTS

I would like to express my deep appreciation and gratitude to my advisor, Dr. Sarah Witherspoon, for all her support, instruction, and encouragement throughout my graduate studies. Her availability and comments during the last stages of the writing of this dissertation are greatly appreciated. It has been a very gratifying experience working with her.

I thank the other members of my advisory committee, Dr. Marcelo Aguiar, Dr. Frank Sottile and Dr. Andreas Klappenecker, for their time, comments on my research, and editorial advice. My special thanks to Dr. Frank Sottile for his constant support, sharing of knowledge and advice since the beginning of my graduate studies.

I thank my fellow graduate students and friends both in Texas A&M and in the University of Massachusetts, for making these last years very enjoyable. Living so many years away from home would have been much harder without them. I also thank my friends in Uruguay for helping me throughout these years with their love and correspondence.

I thank the Department of Mathematics of Texas A&M and its staff, for their help, support and hospitality.

I thank Dr. Walter R. Ferrer from the Centro de Matemática at the Universidad de la República, Uruguay, for his advice that led me to pursue a Ph.D. in mathematics and for his constant support ever since. My gratitude to the Centro de Matemática and Facultad de Ciencias, of the Universidad de la República, for welcoming back at the end of my graduate studies. My special thanks to Sandra Fleitas, who helped me deal with the heavy paperwork involved in this return.

Finally, I would like to thank my parents and sisters for their love, patience and support.

TABLE OF CONTENTS

CHAPTER		Page
I	INTRODUCTION AND PRELIMINARIES	1
	1. Hopf algebras	2
	2. Modules, comodules and Yetter-Drinfel'd modules	8
	3. Radford's construction	13
	4. Some general results	15
II	TWO-PARAMETER QUANTUM GROUPS	18
	1. Definition of restricted quantum groups	18
	2. Factorization of simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules	21
III	COMPUTATIONAL RESULTS	33
	1. G -algebras	33
	2. The code	36
	3. Computational results and conjectures	44
IV	POINTED HOPF ALGEBRAS OF RANK ONE	47
	1. Pointed Hopf algebras of rank one of nilpotent type	47
	2. Factorization of simple $D(H_{\mathcal{D}})$ -modules	48
V	CONCLUSION	60
	REFERENCES	61
	VITA	64

CHAPTER I

INTRODUCTION AND PRELIMINARIES

I study the simple modules of two types of pointed Hopf algebras: restricted two-parameter quantum groups and the Drinfel'd double of rank one pointed Hopf algebras of nilpotent type. The main tool I use is a construction introduced by Radford [19] where the simple modules for the Drinfel'd double of a Hopf algebra are parametrized by group-like elements of the Drinfel'd double.

The dissertation is organized as follows. In this chapter I give the definitions and notations that I will use and I present Radford's construction for simple modules for the Drinfel'd double of certain Hopf algebras. In Chapter II, I define the two-parameter quantum groups and present a theorem on factorization of their simple modules. In Chapter III, I show the code used to construct these modules using the computer algebra system `SINGULAR::PLURAL` and I formulate conjectures about their bases and dimensions based on the computational results. In Chapter IV, I present the rank one pointed Hopf algebras of nilpotent type defined by Krop and Radford in [15], and give a theorem about the reducibility of the tensor product of two simple modules for their Drinfel'd doubles.

In what follows \mathbb{K} is a field of characteristic 0. All vector spaces and tensor products are over \mathbb{K} . A *map* between vector spaces means a linear transformation. For a map $T : V \rightarrow W$ between vector spaces V and W , I will denote the *dual* of T by T^* ; that is $T^* : W^* \rightarrow V^*$ and $T(f)(v) = f(T(v))$ for all $f \in W^*$ and $v \in V$. For vector spaces V and W , the *twist map* $\tau : V \otimes W \rightarrow W \otimes V$ is given by $\tau(v \otimes w) = w \otimes v$.

The journal model is Journal of Algebra.

1. Hopf algebras

I give a brief introduction to Hopf algebras, summarizing the first chapter of [18].

Definition I.1. An *algebra* is a triple (A, m, u) where A is a vector space and

$$m : A \otimes A \rightarrow A \text{ and } u : \mathbb{K} \rightarrow A$$

are maps so that the following diagrams commute:

$$\begin{array}{ccc}
 A \otimes A \otimes A & \xrightarrow{m \otimes id} & A \otimes A \\
 \downarrow id \otimes m & & \downarrow m \\
 A \otimes A & \xrightarrow{m} & A
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & A \otimes A & & \\
 & u \otimes id & \nearrow & id \otimes u & \\
 \mathbb{K} \otimes A & & A \otimes A & & A \otimes \mathbb{K} \\
 \searrow \cong & & \downarrow m & & \nearrow \cong \\
 & & A & &
 \end{array}
 .$$

These are the diagrams of *associativity* and *unit* respectively. The map m is called *multiplication* and u is the *unit*.

Write $m(a \otimes b) = ab$ and $u(1_{\mathbb{K}}) = 1_A$. With this notation, the commutativity of the diagrams means $(ab)c = a(bc)$ and $a1_A = 1_Aa = a$, $\forall a, b, c \in A$. When there is no place for confusion I will say the algebra A instead of (A, m, u) .

Now I dualize the notions just defined to define coalgebras.

Definition I.2. A *coalgebra* is a triple (C, Δ, ε) where C is a vector space and

$$\Delta : C \rightarrow C \otimes C \text{ and } \varepsilon : C \rightarrow \mathbb{K}$$

are maps so that the following diagrams commute:

$$\begin{array}{ccc}
C & \xrightarrow{\Delta} & C \otimes C \\
\Delta \downarrow & & \downarrow \Delta \otimes id \\
C \otimes C & \xrightarrow{id \otimes \Delta} & C \otimes C \otimes C
\end{array}
\qquad
\begin{array}{ccccc}
& & C & & \\
& \nearrow \mathbb{R} & \downarrow \Delta & \nwarrow \mathbb{R} & \\
\mathbb{K} \otimes C & & C \otimes C & & C \otimes \mathbb{K} \\
& \nwarrow \varepsilon \otimes id & & \nearrow id \otimes \varepsilon & \\
& & C \otimes C & &
\end{array}$$

These are the *coassociativity* and *counit* diagrams respectively. The map Δ is called *comultiplication* and ε is the *counit*.

The following notation was introduced by Heyneman and Sweedler.

Notation. The *sigma notation* for Δ is given as follows: for any $c \in C$, write

$$\Delta(x) = \sum x_{(1)} \otimes x_{(2)}.$$

The subscripts (1) and (2) are symbolic and do not indicate particular elements of C .

With this notation the coassociativity diagram translates as

$$\sum x_{(1)(1)} \otimes x_{(1)(2)} \otimes x_{(2)} = \sum x_{(1)} \otimes x_{(2)(1)} \otimes x_{(2)(2)}.$$

This element is denoted by

$$\Delta_2(x) = \sum x_{(1)} \otimes x_{(2)} \otimes x_{(3)}.$$

Iterating this process, applying coassociativity $n - 1$ times, gives

$$\Delta_{n-1}(x) = \sum x_{(1)} \otimes \cdots \otimes x_{(n)}.$$

The counit diagram says that, for all $c \in C$

$$\sum \varepsilon(c_{(1)})c_{(2)} = c = \sum \varepsilon(c_{(2)})c_{(1)}.$$

Definition I.3. Let (C, Δ, ε) be a coalgebra and I a subspace of C .

1. I is a *left coideal* of C if $\Delta(I) \subset C \otimes I$.
2. I is a *right coideal* of C if $\Delta(I) \subset I \otimes C$.
3. I is a *coideal* of C if $\Delta(I) \subset I \otimes C + C \otimes I$ and $\varepsilon(I) = 0$.

If I is a coideal of (C, Δ, ε) , then C/I is a coalgebra with comultiplication and counit induced from Δ and ε respectively.

Example I.4. If (A, m, u) is a finite-dimensional algebra then its dual, A^* , is a coalgebra with $\Delta = m^*$ and $\varepsilon = u^*$. Explicitly, if $f \in A^*$, then $\Delta(f)(a \otimes b) = \sum f_{(1)}(a)f_{(2)}(b) = f(ab)$ for all a and b in A , and $\varepsilon(f) = f(1_A)$.

If (C, Δ, ε) is a coalgebra, then C^* is an algebra with $m = \Delta^*$ and $u = \varepsilon^*$. That is, for f and g in C^* , $(fg)(c) = \sum f(c_{(1)})g(c_{(2)})$ for all $c \in C$ and $1_{C^*} = \varepsilon$.

Definition I.5. A *bialgebra* is a quintuple $(B, m, u, \Delta, \varepsilon)$ where (B, m, u) is an algebra, (B, Δ, ε) is a coalgebra, and the maps Δ and ε are algebra morphisms (or equivalently, m and u are coalgebra morphisms).

Example I.6. If $(B, m, u, \Delta, \varepsilon)$ is a bialgebra, then so are $B^{\text{op}} = (B, m^{\text{op}}, u, \Delta, \varepsilon)$ and $B^{\text{coop}} = (B, m, u, \Delta^{\text{op}}, \varepsilon)$, with $m^{\text{op}} = m \circ \tau$ and $\Delta^{\text{op}} = \tau \circ \Delta$. If $m^{\text{op}} = m$ then B is *commutative*, and if $\Delta^{\text{op}} = \Delta$ it is *cocommutative*.

Definition I.7. Let (A, m, u) be an algebra and (C, Δ, ε) a coalgebra. Then $\text{Hom}_{\mathbb{K}}(C, A)$, the set of linear maps from C to A , is an algebra with the *convolution product*

$$f * g := m \circ (f \otimes g) \circ \Delta$$

for all $f, g \in \text{Hom}_{\mathbb{K}}(C, A)$; *i.e.*

$$(f * g)(x) = \sum f(x_{(1)})g(x_{(2)}), \forall x \in C.$$

The unit element in $\text{Hom}_{\mathbb{K}}(C, A)$ is $u\varepsilon$.

From now on, when I say the algebra $\text{Hom}_{\mathbb{K}}(C, A)$, I mean $(\text{Hom}_{\mathbb{K}}(C, A), *, u \circ \varepsilon)$. In particular, if $(B, m, u, \Delta, \varepsilon)$ is a bialgebra, then $\text{Hom}_{\mathbb{K}}(B, B)$ is an algebra with the structure just described. The map id_B is invertible in $\text{Hom}_{\mathbb{K}}(B, B)$ if and only if there exists a map $S : B \rightarrow B$ such that $S * \text{id}_B = \text{id}_B * S = u \circ \varepsilon$. In other words,

$$\sum S(x_{(1)})x_{(2)} = \sum x_{(1)}S(x_{(2)}) = \varepsilon(x)1_B, \forall x \in B.$$

Such a map S is called an *antipode* in B . If an antipode exists in $(B, m, u, \Delta, \varepsilon)$, it is unique.

Definition I.8. A *Hopf algebra* is a sextuple $(H, m, u, \Delta, \varepsilon, S)$ where $(H, m, u, \Delta, \varepsilon)$ is a bialgebra and $S : H \rightarrow H$ is an antipode in H .

A subspace I of H is a *Hopf ideal* of H , if it is both an ideal and a coideal and $S(I) \subseteq I$. If I is a Hopf ideal of H , then H/I is a Hopf algebra with the structure induced from H .

Example I.9. If (G, \cdot, e) is a group, let $\mathbb{K}G$ be the vector space with basis G . Then $\mathbb{K}G$ is a Hopf algebra with the operations defined by

$$m(g \otimes g') = g \cdot g' \text{ and } u(1) = e, \forall g, g' \in G,$$

$$\Delta(g) = g \otimes g, \varepsilon(g) = 1, \text{ and } S(g) = g^{-1}, \forall g \in G.$$

The algebra $\mathbb{K}G$ is called *the group algebra of G* .

For any coalgebra C , an element $c \in C$ is called *group-like* if

$$\Delta(c) = c \otimes c \text{ and } \varepsilon(c) = 1.$$

Denote by $G(C)$ the set of group-like elements of C . Then $\mathbb{K}G(C)$ is a subcoalgebra of C .

Example I.10. Let \mathfrak{g} be a Lie algebra over \mathbb{K} . The *universal enveloping algebra* $U(\mathfrak{g})$ is the quotient of the tensor algebra $T(\mathfrak{g})$ by the ideal generated by the relations $h \otimes g - g \otimes h - [h, g]$ for all h, g in \mathfrak{g} . Then $U(\mathfrak{g})$ is a Hopf algebra with:

$$\Delta(h) = h \otimes 1 + 1 \otimes h, \quad \varepsilon(h) = 0, \quad \text{and } S(h) = -h, \quad \forall h \in \mathfrak{g}.$$

Example I.11. If H is a finite-dimensional Hopf algebra with antipode S , then H^* with the structures described in I.4 is a Hopf algebra with antipode S^* .

Example I.12. In [21] Taft constructed a family of finite-dimensional non-commutative, non-cocommutative Hopf algebras: let $\ell \in \mathbb{Z}_{>0}$, and θ a primitive ℓ th root of unity. The Taft algebra T_θ is generated as an algebra by elements x and a , subject to the relations:

$$x^\ell = 0, \quad a^\ell = 1, \quad ax = \theta xa.$$

The coalgebra structure and the antipode are determined by:

$$\Delta(a) = a \otimes a, \quad \varepsilon(a) = 1, \quad S(a) = a^{-1} = a^{\ell-1},$$

$$\Delta(x) = x \otimes a + 1 \otimes x. \quad \varepsilon(x) = 0, \quad S(x) = -xa^{-1}.$$

The set $\{a^i x^j : 0 \leq i, j < \ell\}$ is a linear basis for T_θ .

The Hopf algebras that I will study are generalizations of the Taft algebras, and they will all be graded Hopf algebras, as defined next.

Definition I.13. A Hopf algebra H is *graded* if $H = \bigoplus_{n=0}^{\infty} H_n$ and

1. H is a graded algebra, *i.e.* $1 \in H_0$ and $H_m H_n \subseteq H_{m+n}$.
2. H is a graded coalgebra, *i.e.* $\Delta(H_n) \subseteq \sum_{i=0}^n H_{n-i} \otimes H_i$ and $\varepsilon(H_n) = 0, \forall n > 0$.
3. $S(H_n) \subseteq H_n, \forall n \geq 0$.

The Taft algebra T_θ is a graded Hopf algebra with $(T_\theta)_n = \mathbb{K}\{a^i x^n : 0 \leq i < \ell\}$ if $n < \ell$, and $(T_\theta)_n = (0)$ for $n \geq \ell$.

Another property of the Taft algebras is that they are pointed Hopf algebras, as defined next.

Definition I.14. A coalgebra is called *simple* if it has no proper subcoalgebras. For a coalgebra C , the *coradical* $C_{(0)}$ of C , is the sum of the simple subcoalgebras of C . If $C_{(0)} = \mathbb{K}G(C)$ (in other words, every simple subcoalgebra of C is one-dimensional), C is *pointed*.

Now I give a definition that will be used in the following chapters. Given any Hopf algebra H and L a subset of H , let

$$L^+ = L \cap \text{Ker } \varepsilon.$$

Note that if L is a subcoalgebra of H , then L^+ is a coideal and hence H/L^+ is a coalgebra. Moreover, let $\langle L^+ \rangle = HL^+H$ be the two-sided ideal generated by L^+ , then $H/\langle L^+ \rangle$ is a bialgebra. I will use this construction in the particular case where $L \subset Z(H)$, the center of H , in which case $\langle L^+ \rangle = HL^+$ and so H/HL^+ is a bialgebra. If in addition $S(L^+) \subset L^+$, then H/HL^+ is a Hopf algebra. A simple calculation shows that if $L = \mathbb{K}J$ with J a subgroup of $G(H)$, the group of group-like elements of H , then

$$L^+ = \mathbb{K}\{g - 1 : g \in J\}.$$

Remark I.15. In [20] H.-J. Schneider strengthened the Nichols-Zoeller theorem and showed that if H is a finite-dimensional Hopf algebra and L is a Hopf subalgebra of H , then $H \simeq H/HL^+ \otimes L$ as right L -modules [20]. In particular

$$\dim(H/HL^+) = \frac{\dim(H)}{\dim(L)}.$$

Definition I.16. For H a finite-dimensional Hopf algebra, let

$$G_C(H) = G(H) \cap Z(H)$$

denote the group of central group-like elements of H and let

$$\overline{H} = H/H(\mathbb{K}G_C(H))^+.$$

Then \overline{H} is a Hopf algebra, and by Remark I.15

$$\dim(\overline{H}) = \frac{\dim(H)}{|G_C(H)|}.$$

2. Modules, comodules and Yetter-Drinfel'd modules

Definition I.17. Let A be an algebra. A *left A -module* is a pair (M, ρ) , where M is a vector space and $\rho : A \otimes M \rightarrow M$ is a map so that the following diagrams commute:

$$\begin{array}{ccc} A \otimes A \otimes M & \xrightarrow{m \otimes id} & A \otimes M, \\ id \otimes \rho \downarrow & & \downarrow \rho \\ A \otimes M & \xrightarrow{\rho} & A \end{array} \quad \begin{array}{ccc} \mathbb{K} \otimes M & \xrightarrow{u \otimes id} & A \otimes M. \\ & \searrow \cong & \downarrow \rho \\ & & M \end{array}$$

A map ρ as above is called an *action*. Write $\rho(a \otimes m) = a \cdot m$. With this notation the diagrams become

$$a \cdot (b \cdot m) = (ab) \cdot m \text{ and } 1_A \cdot m = m, \forall a, b \in A, m \in M.$$

There is an analogous definition of *right modules*; since all the modules I will consider will be left modules, I will say *module* for left module.

If (M, ρ_M) and (N, ρ_N) are A -modules, a map $f : M \rightarrow N$ is a *morphism of modules* if $f(a \cdot m) = a \cdot f(m)$, $\forall m \in M, a \in A$.

Definition I.18. Let (M, ρ_M) be an A -module and N a subspace of M ; N is an

A -submodule of M if $A \cdot N \subset N$. An A -module M is *simple* if its only submodules are 0 and M . A modules is *completely reducible* if it is the direct sum of its simple submodules.

Dualizing the previous definitions we get the analogous notions for coalgebras.

Definition I.19. Let C be a coalgebra. A *right C -comodule* is a pair (M, δ) where M is a vector space and $\delta : M \rightarrow M \otimes C$ is a map so that the following diagrams commute:

$$\begin{array}{ccc} M & \xrightarrow{\delta} & M \otimes C \\ \delta \downarrow & & \downarrow id \otimes \Delta \\ M \otimes C & \xrightarrow{\delta \otimes id} & M \otimes C \otimes C \end{array} , \quad \begin{array}{ccc} M & \xrightarrow{\delta} & M \otimes C \\ \parallel \searrow & & \downarrow id \otimes \varepsilon \\ M & & M \otimes \mathbb{K} \end{array} .$$

The map δ is called *coaction*. Write

$$\delta(m) = \sum m_{(0)} \otimes m_{(1)},$$

where $m_{(0)} \in M$, $m_{(1)} \in C$.

Remark I.20. With this notation, the diagrams above translate as

$$\sum m_{(0)(0)} \otimes m_{(0)(1)} \otimes m_{(1)} = \sum m_{(0)} \otimes m_{(1)(1)} \otimes m_{(1)(2)} \quad (\text{I.1})$$

and

$$\sum m_{(0)} \varepsilon(m_{(1)}) = m,$$

for all $m \in M$. The element of the equation (I.1) will be denoted $\sum m_{(0)} \otimes m_{(1)} \otimes m_{(2)}$.

Definition I.21. Given (M, δ) and (N, η) two C -comodules, a map $f : M \rightarrow N$ is a

comodule morphism if the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \delta_M \downarrow & & \downarrow \delta_N \\ M \otimes C & \xrightarrow{f \otimes id} & N \otimes C \end{array} .$$

That is, if $\sum f(m_{(0)}) \otimes m_{(1)} = \sum f(m)_{(0)} \otimes f(m)_{(1)}, \forall m \in M$.

There is an analogous definition of *left comodule*; since all the comodules will be right comodules, I will say *comodule* for right comodule.

Definition I.22. Let (M, δ) be a C -comodule and N a subspace of M ; N is a C -subcomodule of M if $\delta(N) \subset N \otimes C$.

Remark I.23. If $(B, m, u, \Delta, \varepsilon)$ is a bialgebra and M and N are B -modules, then $M \otimes N$ is also B -module with action given by

$$b \cdot (m \otimes n) = \sum b_{(1)} \cdot m \otimes b_{(2)} \cdot n, \forall b \in B, m \in M, n \in N.$$

If M and N are B -comodules then $M \otimes N$ is a B -comodule with coaction

$$\delta(m \otimes n) = \sum m_{(0)} \otimes n_{(0)} \otimes m_{(1)} n_{(1)}, \forall m \in M, n \in N.$$

Definition I.24. Let H be a finite-dimensional Hopf algebra over \mathbb{K} with antipode S . The *Drinfel'd double* of H , $D(H)$, is

$$D(H) = (H^*)^{\text{coop}} \otimes H$$

as a coalgebra. The algebra structure is given by

$$(g \otimes h)(f \otimes k) = \sum g(h_{(1)} \rightharpoonup f \leftarrow S^{-1}(h_{(3)})) \otimes h_{(2)}k,$$

for all $g, f \in H^*$ and $h, k \in H$; where $(a \rightharpoonup f)(b) = f(ba)$ and $(f \leftarrow a)(b) = f(ab)$,

for all $a, b \in H$ and $f \in H^*$.

This construction is due to Drinfel'd [10] where he showed that if H is a finite-dimensional Hopf algebra, then $D(H)$ is a Hopf algebra. Furthermore, if M and N are $D(H)$ -modules, then

$$M \otimes N \simeq N \otimes M.$$

Explicitly, if $\{h_i\}$ is a basis for H and $\{h_i^*\}$ is the corresponding dual basis of H^* , let

$$R = \sum_i (\varepsilon_H \otimes h_i) \otimes (h_i^* \otimes 1_H) \in D(H) \otimes D(H).$$

Then $M \otimes N \simeq N \otimes M$ via $m \otimes n \mapsto R^{-1}(n \otimes m)$. Drinfel'd doubles are examples of *quasitriangular bialgebras*, which are bialgebras B equipped with invertible elements $R \in B \otimes B$, satisfying certain conditions, and for which the symmetry of tensor products of modules is realized via R^{-1} .

If M is a $D(H)$ -module, then it is both an H -module and an $(H^*)^{\text{coop}}$ -module. The action of H^* gives rise to an H -comodule structure on M such that if $\delta(m) = \sum m_{(0)} \otimes m_{(1)}$ then $f \cdot m = \sum \langle f, m_{(1)} \rangle m_{(0)}$ for all $f \in H^*$.

Definition I.25. For any bialgebra H , a *left-right Yetter-Drinfel'd module* is a \mathbb{K} -vector space M which is both a left H -module and a right H -comodule, and satisfies the compatibility condition

$$\sum h_{(1)} \cdot m_{(0)} \otimes h_{(2)} m_{(1)} = \sum (h_{(2)} \cdot m)_{(0)} \otimes (h_{(2)} \cdot m)_{(1)} h_{(1)}.$$

The category of left-right Yetter-Drinfel'd modules over a bialgebra H will be denoted by ${}_H\mathcal{YD}^H$.

Proposition I.26 (Majid [17]). *Let H be a finite dimensional Hopf algebra. Then $D(H)$ -modules are left-right Yetter-Drinfel'd modules and conversely. Explicitly, if M is a left-right Yetter-Drinfel'd module, then it is a $D(H)$ -module with the same*

action of H and the action of H^* given by

$$f \cdot m = \sum f(m_{(1)})m_{(0)}, \quad (\text{I.2})$$

for all f in H^* and m in M .

Remark I.27. If $M, N \in {}_H\mathcal{YD}^H$, by the last proposition M and N are $D(H)$ -modules. Since $D(H)$ is a bialgebra, by Remark I.23 $M \otimes N$ is also a $D(H)$ -module and hence a Yetter-Drinfel'd module over H . The Yetter-Drinfel'd structure is given by

$$h \cdot (m \otimes n) = \sum h_{(1)} \cdot m \otimes h_{(2)} \cdot n$$

and

$$\delta(m \otimes n) = \sum m_{(0)} \otimes n_{(0)} \otimes n_{(1)}m_{(1)}.$$

An alternative definition of the Drinfel'd double is $D'(H) = H \otimes (H^*)^{\text{coop}}$ as coalgebras, and multiplication given by

$$(k \otimes f)(h \otimes g) = \sum kf_{(1)}(S^{-1}(h_{(1)}))f_{(3)}(h_{(3)})h_{(2)} \otimes f_{(2)}g,$$

where $(\Delta^{\text{op}} \otimes \text{id})\Delta^{\text{op}}(f) = \sum f_{(1)} \otimes f_{(2)} \otimes f_{(3)}$. I will need both definitions of the Drinfel'd double since two of the papers I will be using [6, 19] use these different definitions. The following lemma gives the relationship between these two definitions of the Drinfel'd double.

Lemma I.28. $D'(H) \simeq D(H^*)^{\text{coop}}$ as Hopf algebras.

Proof. As $H^{**} \simeq H$, we have $D(H^*) \cong H^{\text{coop}} \otimes H^*$, with multiplication

$$\begin{aligned} (k \otimes f)(h \otimes g) &= \sum k(f_{(1)} \rightharpoonup h \leftarrow (S^*)^{-1}(f_{(3)})) \otimes f_{(2)}g \\ &= \sum k(f_{(1)}(h_{(2)})h_{(1)} \leftarrow (f_{(3)} \circ S^{-1})) \otimes f_{(2)}g \\ &= \sum kf_{(1)}(h_{(3)})(f_{(3)}(S^{-1}(h_{(1)}))h_{(2)} \otimes f_{(2)}g, \end{aligned}$$

where $(\Delta^{\text{op}} \otimes \text{id})\Delta^{\text{op}}(f) = \sum f_{(3)} \otimes f_{(2)} \otimes f_{(1)}$. So $D'(H) \simeq D(H^*)$ as algebras. As coalgebras $D(H^*) \simeq H^{\text{coop}} \otimes H^* = (H \otimes (H^*)^{\text{coop}})^{\text{coop}} = D'(H)^{\text{coop}}$. \square

3. Radford's construction

In this section I describe results from [19]. Although Radford's results are more general, I will only write them for \mathbb{K} an algebraically closed field of characteristic 0. This is the main tool I will use to study representations of Drinfel'd doubles. For algebras A and B , the set of algebra maps from A to B will be denoted by $\text{Alg}(A, B)$. It is not hard to see that if H is a finite dimensional algebra, then $\text{Alg}(H, \mathbb{K}) = G(H^*)$, the set of group-like elements of H^* .

Lemma I.29 (Radford [19]). *Let H be a bialgebra over \mathbb{K} and suppose H^{op} is a Hopf algebra with antipode \bar{S} . If $\beta \in \text{Alg}(H, \mathbb{K})$, then $H_\beta = (H, \bullet_\beta, \Delta) \in {}_H\mathcal{YD}^H$, where*

$$h \bullet_\beta a = \sum \beta(h_{(3)})h_{(2)}a\bar{S}(h_{(1)}), \quad (\text{I.3})$$

for all h, a in H .

If $\beta : H \rightarrow \mathbb{K}$ is an algebra map and N is a right coideal of H , then the H -submodule of H_β generated by N , $H \bullet_\beta N$, is a Yetter-Drinfel'd H -submodule of H_β . If $g \in G(H)$, then $\mathbb{K}g$ is a right coideal and $H \bullet_\beta \mathbb{K}g = H \bullet_\beta g$ is a Yetter-Drinfel'd submodule of H_β . For M a Yetter-Drinfel'd module over H , $[M]$ will denote the isomorphism class of M .

Proposition I.30 (Radford [19]). *Let $H = \bigoplus_{n=0}^{\infty} H_n$ be a graded Hopf algebra over \mathbb{K} . Suppose that $H_0 = \mathbb{K}G$ where G is a finite abelian group and $H_n = H_{n+1} = \dots = (0)$ for some $n > 0$. Then*

$$(\beta, g) \mapsto [H \bullet_\beta g]$$

is a bijective correspondence between the Cartesian product of sets $\text{Alg}(H, \mathbb{K}) \times G$ and the set of isomorphism classes of simple Yetter-Drinfel'd H -modules.

Let $H = \bigoplus_{n=0}^{\infty} H_n$ be a graded coalgebra and $h = h_0 + \cdots + h_n$ a group-like element of H with $h_i \in H_i$ and $h_n \neq 0$. The coalgebra grading implies that $\Delta(h) \in \sum_{m=0}^n (\sum_{i=0}^m H_{m-i} \otimes H_i)$, but since h is a group-like element $\Delta(h) = h \otimes h = \sum_{i,j=0}^n h_i \otimes h_j \notin \sum_{m=0}^n (\sum_{i=0}^m H_{m-i} \otimes H_i)$ unless $n = 0$. Hence $G(H) = G(H_0)$. In the case where $H_0 = \mathbb{K}G$ we have $G(H) = G(H_0) = G(\mathbb{K}G) = G$, the last equality holding since distinct group-like elements are linearly independent. If H is as in the last proposition is also finite dimensional, then $\text{Alg}(H, \mathbb{K}) \times G = G(H^*) \times G(H)$.

Remark I.31. Let $H = \bigoplus_{n=0}^{\infty} H_n$ be a graded Hopf algebra with $H_m = H_{m+1} = \cdots = (0)$ for some $m > 0$ and $H_0 = \mathbb{K}G(H)$ where $G(H) = G$ is a finite group. If $\beta : H \rightarrow \mathbb{K}$ is an algebra map and $i > 0$, since $H_i^m = (0)$ we have that $\beta|_{H_i} = 0$. Then β is determined by its restriction to $H_0 = \mathbb{K}G$. Since G is a finite group, $1 = \beta(g^{|G|}) = \beta(g)^{|G|}$ and so $\beta(g) \neq 0$ for all $g \in G$. Let

$$\widehat{G} = \text{Hom}(G, \mathbb{K}^\times), \quad (\text{I.4})$$

the set of group homomorphisms from G to $\mathbb{K}^\times = \mathbb{K} - \{0\}$. Then, to give an algebra map $\beta : H \rightarrow \mathbb{K}$, is equivalent to giving a map in \widehat{G} ; when no confusion arises, the corresponding map in \widehat{G} will also be called β .

Example I.32. Let $H = H_0 = \mathbb{K}G$ with G a finite abelian group. If $\beta \in \widehat{G}$ and $g, h \in G$, then $h \bullet_\beta g = \beta(h)g$ and so $H \bullet_\beta g = \mathbb{K}g$. In this case $D(H) = \mathbb{K}\widehat{G} \otimes \mathbb{K}G$ with multiplication given by $(\alpha \otimes h)(\beta \otimes g) = \alpha\beta \otimes hg$. A pair $(\beta, g) \in \widehat{G} \times G$ is then a character of $G \times \widehat{G}$ via $(\beta, g)((h, \alpha)) = \beta(h)\alpha(g)$, $\forall h \in G$ and $\alpha \in \widehat{G}$. The simple Yetter-Drinfel'd module $H \bullet_\beta g$ is then a $D(H)$ -module with action

$$(\alpha \otimes h) \cdot g = \alpha(g)\beta(h)g = (\beta, g)((h, \alpha))g.$$

4. Some general results

I first start by presenting some general results on the tensor product of Yetter-Drinfel'd modules. Throughout this section $H = \bigoplus_{n=0}^{\infty} H_n$ is a graded Hopf algebra over an algebraically closed field \mathbb{K} , $H_0 = \mathbb{K}G$ where G is a finite abelian group and $H_m = H_{m+1} = \cdots = (0)$ for some $m > 0$.

Proposition I.33. *Let $\beta, \beta' \in \text{Alg}(H, \mathbb{K})$ and $g, g' \in G(H)$. If $H_{\bullet\beta}g \otimes H_{\bullet\beta'}g'$ is a simple Yetter-Drinfel'd module, then*

$$H_{\bullet\beta}g \otimes H_{\bullet\beta'}g' \simeq H_{\bullet\beta*\beta'}gg'$$

Proof. Since $H_{\bullet\beta}g \otimes H_{\bullet\beta'}g'$ is a simple Yetter-Drinfel'd module, by Proposition I.30, there exist unique $\beta'' \in \text{Alg}(H, \mathbb{K})$ and $g'' \in G(H)$ such that

$$H_{\bullet\beta}g \otimes H_{\bullet\beta'}g' \simeq H_{\bullet\beta''}g''$$

as Yetter-Drinfel'd modules. Let $\Phi : H_{\bullet\beta}g \otimes H_{\bullet\beta'}g' \rightarrow H_{\bullet\beta''}g''$ be such an isomorphism. Since Φ is a comodule map, we have

$$\begin{aligned} (\Phi \otimes \text{id}) \circ \delta(g \otimes g') &= \delta \circ \Phi(g \otimes g') \quad \Rightarrow \\ (\Phi \otimes \text{id}) \left(\sum g_{(0)} \otimes g'_{(0)} \otimes g'_{(1)}g_{(1)} \right) &= \Delta(\Phi(g \otimes g')). \end{aligned}$$

Then

$$\Phi(g \otimes g') \otimes g'g = \Delta(\Phi(g \otimes g')). \quad (\text{I.5})$$

This implies that $\mathbb{K}\Phi(g \otimes g')$ is a (simple) right coideal of $H_{\bullet\beta''}g''$. In [19] it was shown that if N is a simple right coideal of H , then the only coideal contained in $H_{\bullet\beta}N$ is N . Therefore $\mathbb{K}\Phi(g \otimes g') = \mathbb{K}g''$ and so $g'' = \lambda\Phi(g \otimes g')$ for some $0 \neq \lambda \in \mathbb{K}$; we may assume that $\lambda = 1$. Applying $\varepsilon \otimes \text{id}$ to both sides of Equation (I.5), we get that

$\Phi(g \otimes g') = \varepsilon(\Phi(g \otimes g'))g'g$. We then have:

$$g'' = \Phi(g \otimes g') = \varepsilon(\Phi(g \otimes g'))g'g.$$

Since distinct group-like elements are linearly independent, this implies that $g'' = g'g$.

Since $(H_i)^m = (0)$ for all $i \geq 1$ we have that $\beta * \beta'(H_i) = (0) = \beta''(H_i)$ for all $i \geq 1$. To show that $\beta'' = \beta * \beta'$ it is then enough to show that they agree on G . Let $h \in G$, then

$$\begin{aligned} \beta''(h)gg' &= h_{\bullet\beta''}gg' = h_{\bullet\beta''}(\Phi(g \otimes g')) = \Phi(h \cdot (g \otimes g')) = \\ &= \Phi(h_{\bullet\beta}g \otimes h_{\bullet\beta'}g') = \Phi(\beta(h)\beta'(h)g \otimes g') = (\beta * \beta')(h)gg', \end{aligned}$$

and so $\beta''(h) = (\beta * \beta')(h)$ for all h in G . \square

If H is any Hopf algebra and $\gamma : H \rightarrow \mathbb{K}$ is an algebra map, then γ has an inverse in $\text{Hom}(H, \mathbb{K})$ given by $\gamma^{-1}(h) = \gamma(S(h))$, since

$$(\gamma * (\gamma \circ S))(h) = \sum \gamma(h_{(1)})\gamma(S(h_{(2)})) = \sum \gamma(h_{(1)}S(h_{(2)})) = \gamma(\epsilon(h)1_H) = \varepsilon(h)1_{\mathbb{K}}.$$

Let $N = \mathbb{K}n$ be a one-dimensional H -module. Then there is an algebra homomorphism $\gamma : H \rightarrow \mathbb{K}$ such that $h \cdot n = \gamma(h)n$ for all $h \in H$. Let \mathbb{K}_γ be \mathbb{K} as a vector space with the action given by $h \cdot 1 = \gamma(h)$, and so $N \simeq \mathbb{K}_\gamma$ as H -modules.

If M is any H -module and $\gamma : H \rightarrow \mathbb{K}$ is an algebra morphism, then the natural vector space isomorphism $M \otimes \mathbb{K}_\gamma \simeq M$ endows M with a new module structure, \cdot' , given by $h \cdot' m = \sum \gamma(h_{(2)})h_{(1)} \cdot m$. I will denote this module by M_γ .

Note that $\mathbb{K}_\gamma \otimes \mathbb{K}_{\gamma^{-1}} \simeq \mathbb{K}_\epsilon$ as H -modules, and therefore for any H -module M ,

$$(M_\gamma)_{\gamma^{-1}} = M_\epsilon = M.$$

Remark I.34. Let H be any Hopf algebra and $\gamma : H \rightarrow \mathbb{K}$ an algebra map. If

M is an H -module and N is a submodule of M , then N_γ is a submodule of M_γ . In particular, M is simple if and only if M_γ is simple.

Let $\text{Soc}(M)$ denote the socle of M , that is, $\text{Soc}(M) = \bigoplus N$, the sum over all simple submodules of M . Then, by the last remark, we have that

$$\text{Soc}(M_\gamma) = (\text{Soc}(M))_\gamma.$$

CHAPTER II

TWO-PARAMETER QUANTUM GROUPS

In 1985 Drinfel'd and Jimbo independently introduced the algebra $U_\theta(\mathfrak{g})$, a one-parameter deformation of the universal enveloping algebra of a semisimple Lie algebra \mathfrak{g} [9, 13]. They were first used to construct solutions to the quantum Yang-Baxter equations and have applications in various areas of mathematics and physics. For θ a root of unity, Lusztig defined the restricted one-parameter quantum group $\mathfrak{u}_\theta(\mathfrak{g})$, a finite-dimensional quotient of $U_\theta(\mathfrak{g})$. In what follows, I give the definitions of the two-parameter versions, $U_{r,s}(\mathfrak{g})$ and $\mathfrak{u}_{r,s}(\mathfrak{g})$ for $\mathfrak{g} = \mathfrak{sl}_n$, the Lie algebra of $n \times n$ matrices of trace 0. These algebras are examples of the algebras constructed by Andruskiewitsch and Schneider in their classification of pointed Hopf algebras with abelian groups of group-like elements. In section 2, I give a theorem about factorization of simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules.

1. Definition of restricted quantum groups

Let \mathbb{K} be an algebraically closed field of characteristic 0 and let $\{\epsilon_1, \dots, \epsilon_n\}$ denote an orthonormal basis of an Euclidean space $E = \mathbb{R}^n$ with an inner product $\langle \cdot, \cdot \rangle$. Let $\alpha_j = \epsilon_j - \epsilon_{j+1}$ ($j = 1, \dots, n-1$). Let $r, s \in \mathbb{K}^\times$ be roots of unity with $r \neq s$ and ℓ be the least common multiple of the orders of r and s . Let θ be a primitive ℓ th root of unity and y and z be nonnegative integers such that $r = \theta^y$ and $s = \theta^z$. Takeuchi defined the following Hopf algebra [22].

Definition II.1. The algebra $U = U_{r,s}(\mathfrak{sl}_n)$ is the unital associative \mathbb{K} -algebra generated by $\{e_j, f_j, \omega_j^{\pm 1}, (\omega'_j)^{\pm 1}, 1 \leq j < n\}$, subject to the following relations.

(R1) The $\omega_i^{\pm 1}, (\omega'_j)^{\pm 1}$ all commute with one another and $\omega_i \omega_i^{-1} = \omega'_j (\omega'_j)^{-1} = 1$,

$$(R2) \quad \omega_i e_j = r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle} e_j \omega_i \quad \text{and} \quad \omega_i f_j = r^{-\langle \epsilon_i, \alpha_j \rangle} s^{-\langle \epsilon_{i+1}, \alpha_j \rangle} f_j \omega_i,$$

$$(R3) \quad \omega'_i e_j = r^{\langle \epsilon_{i+1}, \alpha_j \rangle} s^{\langle \epsilon_i, \alpha_j \rangle} e_j \omega'_i \quad \text{and} \quad \omega'_i f_j = r^{-\langle \epsilon_{i+1}, \alpha_j \rangle} s^{-\langle \epsilon_i, \alpha_j \rangle} f_j \omega'_i,$$

$$(R4) \quad [e_i, f_j] = \frac{\delta_{i,j}}{r-s} (\omega_i - \omega'_i).$$

$$(R5) \quad [e_i, e_j] = [f_i, f_j] = 0 \quad \text{if} \quad |i-j| > 1,$$

$$(R6) \quad e_i^2 e_{i+1} - (r+s) e_i e_{i+1} e_i + r s e_{i+1} e_i^2 = 0, \\ e_i e_{i+1}^2 - (r+s) e_{i+1} e_i e_{i+1} + r s e_{i+1}^2 e_i = 0,$$

$$(R7) \quad f_i^2 f_{i+1} - (r^{-1} + s^{-1}) f_i f_{i+1} f_i + r^{-1} s^{-1} f_{i+1} f_i^2 = 0, \\ f_i f_{i+1}^2 - (r^{-1} + s^{-1}) f_{i+1} f_i f_{i+1} + r^{-1} s^{-1} f_{i+1}^2 f_i = 0,$$

for all $1 \leq i, j < n$.

The following coproduct, counit, and antipode give U the structure of a Hopf algebra:

$$\begin{aligned} \Delta(e_i) &= e_i \otimes 1 + \omega_i \otimes e_i, & \Delta(f_i) &= 1 \otimes f_i + f_i \otimes \omega'_i, \\ \epsilon(e_i) &= 0, & \epsilon(f_i) &= 0, \\ S(e_i) &= -\omega_i^{-1} e_i, & S(f_i) &= -f_i (\omega'_i)^{-1}, \end{aligned}$$

and ω_i, ω'_i are group-like, for all $1 \leq i < n$.

Let U^0 be the group algebra generated by all $\omega_i^{\pm 1}, (\omega'_i)^{\pm 1}$ and let U^+ (respectively, U^-) be the subalgebra of U generated by all e_i (respectively, f_i). Let

$$\mathcal{E}_{j,j} = e_j \quad \text{and} \quad \mathcal{E}_{i,j} = e_i \mathcal{E}_{i-1,j} - r^{-1} \mathcal{E}_{i-1,j} e_i \quad (i > j),$$

$$\mathcal{F}_{j,j} = f_j \quad \text{and} \quad \mathcal{F}_{i,j} = f_i \mathcal{F}_{i-1,j} - s \mathcal{F}_{i-1,j} f_i \quad (i > j).$$

The algebra U has a triangular decomposition $U \cong U^- \otimes U^0 \otimes U^+$ (as vector spaces), and the subalgebras U^+, U^- respectively have monomial Poincaré-Birkhoff-Witt

(PBW) bases [14, 4]

$$\mathcal{E} := \{\mathcal{E}_{i_1, j_1} \mathcal{E}_{i_2, j_2} \cdots \mathcal{E}_{i_p, j_p} \mid (i_1, j_1) \leq (i_2, j_2) \leq \cdots \leq (i_p, j_p) \text{ lexicographically}\}, \quad (\text{II.1})$$

$$\mathcal{F} := \{\mathcal{F}_{i_1, j_1} \mathcal{F}_{i_2, j_2} \cdots \mathcal{F}_{i_p, j_p} \mid (i_1, j_1) \leq (i_2, j_2) \leq \cdots \leq (i_p, j_p) \text{ lexicographically}\}. \quad (\text{II.2})$$

It is shown in [6] that all $\mathcal{E}_{i,j}^\ell$, $\mathcal{F}_{i,j}^\ell$, $\omega_i^\ell - 1$, and $(\omega'_i)^\ell - 1$ ($1 \leq j \leq i < n$) are central in $U_{r,s}(\mathfrak{sl}_n)$. The ideal I_n generated by these elements is a Hopf ideal [6, Thm. 2.17], and so the quotient

$$\mathfrak{u} = \mathfrak{u}_{r,s}(\mathfrak{sl}_n) = U_{r,s}(\mathfrak{sl}_n)/I_n \quad (\text{II.3})$$

is a Hopf algebra, called the *restricted two-parameter quantum group*. Examination of the PBW-bases (II.1) and (II.2) shows that \mathfrak{u} is finite-dimensional and Benkart and Witherspoon showed that \mathfrak{u} is pointed [6, Prop. 3.2].

Let \mathcal{E}_ℓ and \mathcal{F}_ℓ denote the sets of monomials in \mathcal{E} and \mathcal{F} respectively, in which each $\mathcal{E}_{i,j}$ or $\mathcal{F}_{i,j}$ appears as a factor at most $\ell - 1$ times. Identifying cosets in \mathfrak{u} with their representatives, we may assume \mathcal{E}_ℓ and \mathcal{F}_ℓ are basis for the subalgebras of \mathfrak{u} generated by the elements e_i and f_i respectively.

Let \mathfrak{b} be the Hopf subalgebra of $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ generated by $\{\omega_i, e_i : 1 \leq i < n\}$, and \mathfrak{b}' the subalgebra generated by $\{\omega'_i, f_i : 1 \leq i < n\}$.

Benkart and Witherspoon showed that, under some conditions on the parameters r and s , $\mathfrak{b}^* \simeq (\mathfrak{b}')^{\text{coop}}$ as Hopf algebras ([6, Lemma 4.1]). This implies that $\mathfrak{b} \simeq ((\mathfrak{b}')^{\text{coop}})^*$; I present the lemma using the dual isomorphism of the original one.

Lemma II.2. [6, Lemma 4.1] *If $\gcd(y^{n-1} - y^{n-2}z + \cdots + (-1)^{n-1}z^{n-1}, \ell) = 1$ and rs^{-1} is a primitive ℓ th root of unity, then $\mathfrak{b} \simeq ((\mathfrak{b}')^{\text{coop}})^*$ as Hopf algebras. Such an*

isomorphism is given by

$$\langle \omega_i, \omega'_j \rangle = r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle} \quad \text{and} \quad \langle \omega_i, f_j \rangle = 0, \quad (\text{II.4})$$

and

$$\langle e_i, f_j^a g \rangle = \delta_{i,j} \delta_{1,a} \quad \forall g \in G(\mathfrak{b}'). \quad (\text{II.5})$$

Proposition II.3. [6, Thm. 4.8] *Assume $r = \theta^y$ and $s = \theta^z$, where θ is a primitive ℓ th root of unity, and*

$$\gcd(y^{n-1} - y^{n-2}z + \cdots + (-1)^{n-1}z^{n-1}, \ell) = 1.$$

Then there is an isomorphism of Hopf algebras $\mathfrak{u}_{r,s}(\mathfrak{sl}_n) \cong D'(\mathfrak{b}) \cong D((\mathfrak{b}')^{\text{coop}})^{\text{coop}}$.

In the special case $r = \theta$, a primitive ℓ th root of unity, and $s = \theta^{-1}$, $\mathfrak{u} = \mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_n)$ is isomorphic to $D((\mathfrak{b}')^{\text{coop}})$ when n and ℓ are relatively prime.

Under the assumption that $\gcd(y^{n-1} - y^{n-2}z + \cdots + (-1)^{n-1}z^{n-1}, \ell) = 1$, by Proposition II.3, $\mathfrak{u}_{r,s}(\mathfrak{sl}_n) = (D((\mathfrak{b}')^{\text{coop}}))^{\text{coop}}$ and so $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules are Yetter-Drinfel'd modules for $(\mathfrak{b}')^{\text{coop}}$ (only the algebra structure of $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ plays a role when studying $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules, hence $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules are $D((\mathfrak{b}')^{\text{coop}})$ -modules). For simplicity I will denote $H = (\mathfrak{b}')^{\text{coop}}$. Then $G = G(H) = \langle \omega'_i : 1 \leq i < n \rangle$ and H is a graded Hopf algebra with $\omega'_i \in H_0$ and $f_i \in H_1$ for all $1 \leq i < n$ and $H_j = (0)$ if $j \geq 2\ell$. Therefore Proposition I.30 applies to H and isomorphism classes of $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules (or simple Yetter-Drinfel'd H -modules) are in one to one correspondence with $\text{Alg}(H, \mathbb{K}) \times G(H)$.

2. Factorization of simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules

In this section I study under what conditions a simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module can be factored as the tensor product of a one-dimensional module and a simple module which is also

a module for $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)} = \mathfrak{u}_{r,s}(\mathfrak{sl}_n)/\mathfrak{u}_{r,s}(\mathfrak{sl}_n)(\mathbb{K}G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)))^+$. Let ℓ , n , y and z be fixed and θ be a primitive ℓ th root of unity. Let A be the $(n-1) \times (n-1)$ matrix

$$A = \begin{pmatrix} y-z & z & 0 & 0 & \cdots & 0 \\ -y & y-z & z & 0 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & -y & y-z & z \\ 0 & \cdots & \cdots & 0 & -y & y-z \end{pmatrix}$$

The determinant of A is $y^{n-1} - y^{n-2}z + \cdots + (-1)^{n-1}z^{n-1}$. Throughout this section, assume that $\gcd(y^{n-1} - y^{n-2}z + \cdots + (-1)^{n-1}z^{n-1}, \ell) = 1$, and so $\det(A)$ is invertible in $\mathbb{Z}/\ell\mathbb{Z}$. I start by describing the set of central group-like elements in $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$. Clearly $G(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)) = \langle \omega_i, \omega'_i : 1 \leq i < n \rangle$.

Proposition II.4. *A group-like element $g = \omega_1^{a_1} \cdots \omega_{n-1}^{a_{n-1}} \omega'_1{}^{b_1} \cdots \omega'_{n-1}{}^{b_{n-1}}$ is central in $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ if and only if*

$$\begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix} = A^{-1}A^t \begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}$$

in $(\mathbb{Z}/\ell\mathbb{Z})^{n-1}$.

Proof. The element g is central in $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ if and only if $ge_k = e_kg$ and $gf_k = f_kg$ for all $k = 1, \dots, n-1$. By the relations (R2) and (R3) of the definition of $U_{r,s}(\mathfrak{sl}_n)$, for all $k = 1, \dots, n-1$ we have that

$$\begin{aligned} ge_k &= \prod_{i=1}^{n-1} (r^{\langle \epsilon_i, \alpha_k \rangle} s^{\langle \epsilon_{i+1}, \alpha_k \rangle})^{a_i} \prod_{j=1}^{n-1} (r^{\langle \epsilon_{j+1}, \alpha_k \rangle} s^{\langle \epsilon_j, \alpha_k \rangle})^{b_j} e_k g \text{ and} \\ gf_k &= \prod_{i=1}^{n-1} (r^{-\langle \epsilon_i, \alpha_k \rangle} s^{-\langle \epsilon_{i+1}, \alpha_k \rangle})^{a_i} \prod_{j=1}^{n-1} (r^{-\langle \epsilon_{j+1}, \alpha_k \rangle} s^{-\langle \epsilon_j, \alpha_k \rangle})^{b_j} f_k g. \end{aligned}$$

Then g is central if and only if

$$\begin{aligned} 1 &= \prod_{i=1}^{n-1} \left(r^{\langle \epsilon_i, \alpha_k \rangle} s^{\langle \epsilon_{i+1}, \alpha_k \rangle} \right)^{a_i} \prod_{j=1}^{n-1} \left(r^{\langle \epsilon_{j+1}, \alpha_k \rangle} s^{\langle \epsilon_j, \alpha_k \rangle} \right)^{b_j} \\ &= s^{a_{k-1} r^{a_k} s^{-a_k} r^{-a_{k+1}} r^{b_{k-1}} r^{-b_k} s^{b_k} s^{-b_{k+1}}}, \forall k = 1, \dots, n-1, \end{aligned}$$

where $a_0 = a_n = 0 = b_0 = b_n$. Since $r = \theta^y$ and $s = \theta^z$, the last equation holds if and only if

$$z a_{k-1} + (y - z) a_k - y a_{k+1} = (-y b_{k-1} + (y - z) b_k + z b_{k+1}) \pmod{\ell}, \quad (\text{II.6})$$

for all $k = 1, \dots, n-1$. The matrix of coefficients of the left hand side of this system of equations is

$$\begin{pmatrix} y-z & -y & 0 & 0 & \cdots & 0 \\ z & y-z & -y & 0 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & z & y-z & -y \\ 0 & \cdots & \cdots & 0 & z & y-z \end{pmatrix} = A^t$$

and the matrix of coefficients of the right hand side is

$$\begin{pmatrix} y-z & z & 0 & 0 & \cdots & 0 \\ -y & y-z & z & 0 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & -y & y-z & z \\ 0 & \cdots & \cdots & 0 & -y & y-z \end{pmatrix} = A.$$

We then have that g is central if and only if

$$A^t \begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix} = A \begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix}$$

in $(\mathbb{Z}/\ell\mathbb{Z})^{n-1}$. □

Example II.5. For $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_n)$ ($y = 1$ and $z = \ell - 1$), the matrix A is symmetric. Therefore, a group-like element $g = \omega_1^{a_1} \cdots \omega_{n-1}^{a_{n-1}} \omega_1^{b_1} \cdots \omega_{n-1}^{b_{n-1}}$ is central if and only if $b_i = a_i$ for all $i = 1, \dots, n-1$.

In general, $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)(\mathbb{K}G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)))^+ = \mathfrak{u}_{r,s}(\mathfrak{sl}_n)\{g - 1 : g \in G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n))\}$. In particular, by the last example, we have that $\mathfrak{u}_{\theta, \theta^{-1}}(\mathbb{K}G_C(\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_n)))^+$ is generated by $\{\omega_i^{-1} - \omega_i' : i = 1, \dots, n-1\}$. This gives $\overline{\mathfrak{u}_{\theta, \theta^{-1}}} \simeq \mathfrak{u}_{\theta}(\mathfrak{sl}_n)$, the one parameter quantum group.

Henceforth r and s are such that rs^{-1} is also a primitive ℓ th root of unity, that is, $\gcd(y - z, \ell) = 1$.

Remark II.6. If $\beta \in G(H^*)$ and $g = \omega_1^{c_1} \cdots \omega_{n-1}^{c_{n-1}} \in G(H)$, by Proposition I.26, the Yetter-Drinfel'd module $H \bullet_{\beta} g$ is also a $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module where the action of $H^* = \mathfrak{b}$ is given by

$$f \cdot h = \sum \langle f, h_{(2)} \rangle h_{(1)},$$

for all h in $H = (\mathfrak{b}')^{coop}$ and f in $H^* = \mathfrak{b}$. In particular,

$$\omega_i \cdot g = \langle \omega_i, g \rangle g = \prod_{j=1}^{n-1} \langle \omega_i, \omega_j' \rangle^{c_j} g = \prod_{j=1}^{n-1} (r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle})^{c_j} g.$$

Proposition II.7. Let $\beta \in G(H^*)$ be defined by $\beta(\omega_i') = \theta^{\beta_i}$ and $g = \omega_1^{c_1} \cdots \omega_{n-1}^{c_{n-1}}$.

The simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module $H \bullet_{\beta} g$ is naturally a $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$ -module if and only if

$$\begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{n-1} \end{pmatrix} = -A^t \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} \quad (\text{II.7})$$

in $(\mathbb{Z}/\ell\mathbb{Z})^{n-1}$.

Proof. $H \bullet_{\beta} g$ is a $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$ -module if and only if $(h-1) \cdot m = 0$ for all h in $G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n))$ and m in $H \bullet_{\beta} g$. If $h \in G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n))$, $h \cdot m = m$ for all m in $H \bullet_{\beta} g$ if and only if $h \cdot g = g$.

Let $h = \omega_1^{a_1} \cdots \omega_{n-1}^{a_{n-1}} \omega_1^{b_1} \cdots \omega_{n-1}^{b_{n-1}} \in G_C(\mathfrak{u}_{r,s}(\mathfrak{sl}_n))$; then by Proposition II.4

$$\begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix} = A^{-1} A^t \begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}.$$

We have

$$\begin{aligned} \omega_1^{b_1} \cdots \omega_{n-1}^{b_{n-1}} \bullet_{\beta} g &= \beta(\omega_1^{b_1} \cdots \omega_{n-1}^{b_{n-1}}) g \\ &= \theta^{b_1 \beta_1 + \cdots + b_{n-1} \beta_{n-1}} g \end{aligned} \quad (\text{II.8})$$

and

$$\begin{aligned} \omega_1^{a_1} \cdots \omega_{n-1}^{a_{n-1}} \cdot g &= \langle \omega_1^{a_1} \cdots \omega_{n-1}^{a_{n-1}}, g \rangle g \\ &= \prod_{i=1}^{n-1} \langle \omega_i, g \rangle^{a_i} g \\ &= \prod_{i=1}^{n-1} \prod_{j=1}^{n-1} (r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle})^{a_i c_j} g \\ &= \prod_{i=1}^{n-1} (r^{-c_{i-1}} (r s^{-1})^{c_i} s^{c_{i+1}})^{a_i} g \\ &= \theta^x g \end{aligned} \quad (\text{II.9})$$

where $c_0 = c_n = 0$ and $x = \sum_{i=1}^{n-1} (-yc_{i-1} + (y-z)c_i + zc_{i+1}) a_i$. From (II.8) and (II.9) we get that

$$h \cdot g = \theta^{x + \sum_{i=1}^{n-1} b_i \beta_i} g$$

for all $\begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}$, where $\begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix} = A^{-1} A^t \begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}$. Now

$$\sum_{i=1}^{n-1} (-yc_{i-1} + (y-z)c_i + zc_{i+1}) a_i + \sum_{i=1}^{n-1} b_i \beta_i = 0 \pmod{\ell}$$

if and only if

$$\begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}^t A \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} = - \begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix}^t \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{n-1} \end{pmatrix} \text{ in } \mathbb{Z}/\ell\mathbb{Z}.$$

We then have that $H_{\bullet\beta}g$ is a $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$ -module, if and only if

$$\begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}^t A \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} = - \begin{pmatrix} a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}^t A (A^{-1})^t \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{n-1} \end{pmatrix},$$

for all (a_1, \dots, a_{n-1}) in $(\mathbb{Z}/\ell\mathbb{Z})^{n-1}$. This occurs if and only if

$$\begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} = - (A^{-1})^t \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{n-1} \end{pmatrix} \text{ in } (\mathbb{Z}/\ell\mathbb{Z})^{n-1}.$$

□

Given $g = (\omega'_1)^{c_1} \dots (\omega'_{n-1})^{c_{n-1}} \in G(H)$, let β_1, \dots, β_n be defined as in Equation

(II.7). I will denote by β_g the algebra map given by $\beta_g(\omega'_i) = \theta^{\beta_i}$.

For any Hopf algebra H , let \mathcal{S}_H denote the set of isomorphism classes of simple H -modules. Then $\mathcal{S}_{\overline{H}}$ can be identified as the subset of \mathcal{S}_H consisting of the H -modules that are naturally \overline{H} -modules. Combining the last proposition with Proposition I.30, we get

Corollary II.8. *The correspondence $G(H) \rightarrow \mathcal{S}_{\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}}$ given by*

$$g \mapsto [H_{\bullet\beta_g}g]$$

is a bijection.

Example II.9. In the $\mathfrak{u}_{\theta,\theta^{-1}}(\mathfrak{sl}_2)$ case, the matrix A is $A = (2)$. Then, the simple $\mathfrak{u}_{\theta,\theta^{-1}}(\mathfrak{sl}_2)$ -modules that are naturally $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -modules, are of the form $H_{\bullet\beta}(\omega')^c$ with $\beta(\omega') = \theta^{-2c}$.

Example II.10. Using the last Proposition in the case $n = 3$, we have that the $\mathfrak{u}_{r,s}(\mathfrak{sl}_3)$ -module $H_{\bullet\beta}(\omega'_1)^{c_1}(\omega'_2)^{c_2}$ is a $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_3)}$ -module if and only if, $\beta(\omega'_1) = \theta^{(z-y)c_1+y c_2}$ and $\beta(\omega'_2) = \theta^{-z c_1+(z-y)c_2}$. In particular, for the $\mathfrak{u}_{\theta,\theta^{-1}}(\mathfrak{sl}_3)$ -modules, the condition is $\beta(\omega'_1) = \theta^{-2c_1+c_2}$ and $\beta(\omega'_2) = \theta^{c_1-2c_2}$.

For an algebra map $\chi : \mathfrak{u}_{r,s}(\mathfrak{sl}_n) \rightarrow \mathbb{K}$, let \mathbb{K}_{χ} be the 1-dimensional $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module given by $h \cdot 1 = \chi(h)1$. Since $e_i^{\ell} = 0 = f_i^{\ell}$ we have that $\chi(e_i) = \chi(f_i) = 0$, and this together with (R4) of the Definition II.1 of $U_{r,s}(\mathfrak{sl}_n)$, gives that $\chi(\omega_i) = \chi(\omega'_i)$. For each $i = 1, \dots, n-1$, since $\omega_i^{\ell} = 1$, $\chi(\omega_i) = \theta^{\chi_i}$ for some $0 \leq \chi_i < \ell$.

Proposition II.11. *For $\chi : \mathfrak{u}_{r,s}(\mathfrak{sl}_n) \rightarrow \mathbb{K}$ an algebra map, we have that $\mathbb{K}_{\chi} \simeq H_{\bullet\chi|_H}g_{\chi}$, where*

$$g_{\chi} = \omega_1'^{d_1} \cdots \omega_{n-1}'^{d_{n-1}}, \text{ with}$$

$$\begin{pmatrix} d_1 \\ \vdots \\ d_{n-1} \end{pmatrix} = A^{-1} \begin{pmatrix} \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix} \text{ in } (\mathbb{Z}/\ell\mathbb{Z})^{n-1}.$$

Proof. Since \mathbb{K}_χ is a simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module, we have that $\mathbb{K}_\chi \simeq H \bullet_\beta g$ for some unique $\beta \in G(H^*)$ and $g \in G(H)$. Let $\phi : \mathbb{K}_\chi \rightarrow H \bullet_\beta g$ be an isomorphism of Yetter-Drinfel'd modules. We may assume that $g = \phi(1)$; then

$$\beta(\omega'_i)g = \omega'_i \bullet_\beta g = \omega'_i \bullet_\beta (\phi(1)) = \phi(\omega'_i \cdot 1) = \phi(\chi(\omega'_i)1) = \chi(\omega'_i)g.$$

Therefore $\beta(\omega'_i) = \chi(\omega'_i)$ and since $\beta(f_i) = 0 = \chi(f_i)$ for all $i = 1, \dots, n-1$, we have $\beta = \chi|_H$.

We have that

$$\begin{aligned} \omega_i \cdot g &= \langle \omega_i, g \rangle g \\ &= \left(\prod_{j=1}^{n-1} \langle \omega_i, \omega'_j \rangle^{d_j} \right) g \\ &= \left(\prod_{j=1}^{n-1} (r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle})^{d_j} \right) g \\ &= r^{-d_{i-1}} (r s^{-1})^{d_i} s^{d_{i+1}} g \\ &= \theta^{y(d_i - d_{i-1}) + z(d_{i+1} - d_i)} g. \end{aligned} \tag{II.10}$$

On the other hand

$$\omega_i \cdot g = \omega_i \cdot \phi(1) = \phi(\omega_i \cdot 1) = \phi(\theta^{X_i} 1) = \theta^{X_i} g. \tag{II.11}$$

By (II.10) and (II.11) we have that

$$-yd_{i-1} + (y-z)d_i + zd_{i+1} = \chi_i \text{ mod } \ell, \quad \forall i = 1, \dots, n-1; \text{ and so}$$

$$A \begin{pmatrix} d_1 \\ \vdots \\ d_{n-1} \end{pmatrix} = \begin{pmatrix} \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix} \text{ in } (\mathbb{Z}/\ell\mathbb{Z})^{n-1}.$$

□

For any Hopf algebra H , let $\mathcal{S}_H^1 = \{[N] \in \mathcal{S}_H : \dim(N) = 1\}$. Combining the last proposition and Proposition I.30 we get

Corollary II.12. *The correspondence $G(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)^*) \rightarrow \mathcal{S}_{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}^1$ given by*

$$\chi \mapsto [H_{\bullet_{\chi|_H}} g_\chi]$$

is a bijection.

Theorem II.13. *The map $\Phi : \overline{\mathcal{S}_{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}} \times \mathcal{S}_{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}^1 \rightarrow \mathcal{S}_{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$ given by*

$$\Phi([M], [N]) = [M \otimes N]$$

is a bijection if and only if $\gcd((y-z)n, \ell) = 1$.

Proof. By the last corollary we have that 1-dimensional simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -modules are of the form $H_{\bullet_{\chi|_H}} g_\chi$ with $\chi \in G(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)^*)$. Also by Corollary II.8, simple $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$ -modules are of the form $H_{\bullet_{\beta_g}} g$ for $g \in G(H)$. Furthermore by Proposition I.33, we have that $H_{\bullet_{\beta_g}} g \otimes H_{\bullet_{\chi|_H}} g_\chi \simeq H_{\bullet_{\beta_g * \chi}} gg_\chi$. Then Φ is a bijection if and only if

$$\Psi : \{(g, \beta_g) : g \in G(H)\} \times \{(g_\chi, \chi) : \chi \in G(\mathfrak{u}_{r,s}(\mathfrak{sl}_n)^*)\} \rightarrow G(H) \times G(H^*)$$

given by $\Psi((g, \beta_g), (g_\chi, \chi)) = (gg_\chi, \beta_g * \chi|_H)$ is a bijection. The latter holds if and only if for all $h = \omega_1^{b_1} \cdots \omega_{n-1}^{b_{n-1}}$ and γ given by $\gamma(\omega'_i) = \theta^{\gamma_i}$, there exist unique

$g = \omega_1'^{c_1} \cdots \omega_{n-1}'^{c_{n-1}}$ and χ with $\chi(w_i) = \chi(\omega_i') = \theta^{\chi_i}$, so that $h = gg_\chi$ and $\gamma = \beta_g * \chi|_H$.

If $\beta_g(\omega_i') = \theta^{\beta_i}$ and $g_\chi = \omega_1'^{d_1} \cdots \omega_{n-1}'^{d_{n-1}}$, then

$$gg_\chi = \omega_1'^{c_1} \cdots \omega_{n-1}'^{c_{n-1}} \omega_1'^{d_1} \cdots \omega_{n-1}'^{d_{n-1}} \text{ and } (\beta_g * \chi|_H)(\omega_i') = \theta^{\beta_i + \chi_i}.$$

Then Ψ is bijective if and only if the system of equations

$$\begin{pmatrix} c_1 + d_1 \\ \vdots \\ c_{n-1} + d_{n-1} \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix}$$

$$\begin{pmatrix} \beta_1 + \chi_1 \\ \vdots \\ \beta_{n-1} + \chi_{n-1} \end{pmatrix} = \begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_{n-1} \end{pmatrix}$$

subject to

$$\begin{pmatrix} d_1 \\ \vdots \\ d_{n-1} \end{pmatrix} = A^{-1} \begin{pmatrix} \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix}$$

$$\begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{n-1} \end{pmatrix} = -A^t \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix}$$

has a unique solution for all $(b_1, \dots, b_{n-1}), (\gamma_1, \dots, \gamma_{n-1})$. The last four vector equa-

tions are equivalent to

$$\begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} + A^{-1} \begin{pmatrix} \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \end{pmatrix}$$

$$-A^t \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \end{pmatrix} + \begin{pmatrix} \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix} = \begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_{n-1} \end{pmatrix}$$

which can be written as

$$\begin{pmatrix} \text{id} & A^{-1} \\ -A^t & \text{id} \end{pmatrix} \begin{pmatrix} c_1 \\ \vdots \\ c_{n-1} \\ \chi_1 \\ \vdots \\ \chi_{n-1} \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_{n-1} \\ \gamma_1 \\ \vdots \\ \gamma_{n-1} \end{pmatrix}.$$

This last system has a unique solution if and only if the matrix

$$M = \begin{pmatrix} \text{id} & A^{-1} \\ -A^t & \text{id} \end{pmatrix}$$

is invertible in $M_{(n-1) \times (n-1)}(\mathbb{Z}/\ell\mathbb{Z})$, or equivalently, if $\gcd(\det(M), \ell) = 1$. By row-reducing M we have that

$$\begin{aligned}
\det \begin{pmatrix} \text{id} & A^{-1} \\ -A^t & \text{id} \end{pmatrix} &= \det \begin{pmatrix} A & \text{id} \\ -A^t & \text{id} \end{pmatrix} \\
&= \det \begin{pmatrix} A + A^t & 0 \\ -A^t & \text{id} \end{pmatrix} \\
&= \det(A + A^t).
\end{aligned}$$

Now

$$\begin{aligned}
A + A^t &= \begin{pmatrix} 2(y-z) & z-y & 0 & 0 & \cdots & 0 \\ z-y & 2(y-z) & z-y & 0 & \cdots & 0 \\ 0 & z-y & 2(y-z) & z-y & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & z-y & 2(y-z) & z-y \\ 0 & \cdots & \cdots & 0 & z-y & 2(y-z) \end{pmatrix} \\
&= (y-z) \begin{pmatrix} 2 & -1 & 0 & 0 & \cdots & 0 \\ -1 & 2 & -1 & 0 & \cdots & 0 \\ 0 & -1 & 2 & -1 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & -1 & 2 & -1 \\ 0 & \cdots & \cdots & 0 & -1 & 2 \end{pmatrix}.
\end{aligned}$$

Therefore $\det(A + A^t) = (y-z)^{n-1}n$. We then have that Φ is a bijection if and only if $\gcd((y-z)n, \ell) = 1$. \square

CHAPTER III

COMPUTATIONAL RESULTS

In this chapter I present how I used the computer algebra system SINGULAR::PLURAL [12] to construct simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_3)$ -modules. These computations were begun as part of a joint project with G. Benkart and S. Witherspoon to understand the information obtained by Radford's method about $\mathfrak{u}_\theta(\mathfrak{sl}_n)$ -modules [5]. To reduce computations, I use Proposition II.13 and construct only the $\mathfrak{u}_{r,s}(\mathfrak{sl}_3)$ -modules that are also modules for the quotient $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_3)}$ via the quotient map; that is, I only look at the cases when $\gcd((y-z)3, \ell) = 1$. According to Example II.10, we only need to construct the modules $H_{\beta}(\omega'_1)^{c_1}(\omega'_2)^{c_2}$ where $\beta(\omega'_1) = \theta^{(z-y)c_1+yc_2}$ and $\beta(\omega'_2) = \theta^{-zc_1+(z-y)c_2}$.

1. **G-algebras**

The system SINGULAR::PLURAL allows us to do computations on G -algebras, which are algebras given by generators and re-writing relations where Gröbner basis computations can be done. I will give the precise definition of G -algebras and show that $H = (\mathfrak{b}')^{\text{coop}}$ is a quotient of a G -algebra. The notion of G -algebras was introduced by Apel in [2] and later refined by Levandovskyy in [16], and is a generalization of commutative polynomial rings.

Let $T = \mathbb{K}\langle x_1, \dots, x_m \rangle$, the associative algebra generated by x_1, \dots, x_m . The *standard monomials* in A , are elements from the set

$$\text{Mon}_S(A) = \{x^\alpha = x_1^{\alpha_1} \cdots x_m^{\alpha_m} : \alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{N}^m\}.$$

A relation $<_A$ on $\text{Mon}_S(A)$ is called a *monomial ordering on* $\text{Mon}_S(A)$ if the following relations hold:

- $<_A$ is a total well-ordering.
- If $x^\alpha <_A x^\beta$ and $x^\gamma \in \text{Mon}_s(A)$, then $x^{\alpha+\gamma} <_A x^{\beta+\gamma}$.

The *degree* of a monomial $x^\alpha = x_1^{\alpha_1} \cdots x_m^{\alpha_m} \in \text{Mon}_s(A)$ is $\deg(x^\alpha) = \alpha_1 + \cdots + \alpha_m$. For an element $0 \neq f \in \mathbb{K}\text{Mon}_s(A)$, the leading monomial of f with respect to $<_A$ will be denoted by $\text{lm}(f)$. An example of a monomial ordering is the *degree lexicographic* order, $<_{\text{dlex}}$ which is defined by $x^\alpha <_{\text{dlex}} x^\beta$ if $\deg(x^\alpha) < \deg(x^\beta)$ or if $\deg(x^\alpha) = \deg(x^\beta)$ and the left-most nonzero entry of $\beta - \alpha$ is positive. With this order we have $x_1 >_{\text{dlex}} x_2 >_{\text{dlex}} \cdots >_{\text{dlex}} x_m$.

Definition III.1. Let \mathbb{K} be a field and A be an algebra given in terms of generators and relations:

$$A = \mathbb{K}\langle x_1, \dots, x_k \mid x_j x_i = C_{ij} x_i x_j + D_{ij}, \forall 1 \leq i < j \leq k \rangle,$$

where the $C_{ij} \in \mathbb{K}^\times$ and $D_{ij} \in \mathbb{K}\text{Mon}_s(A)$. A is a G-algebra if the following conditions hold:

- There is a monomial well-ordering on $\text{Mon}_s(A)$, $<_A$, such that $\text{lm}(D_{ij}) <_A x_i x_j$ for all $1 \leq i < j \leq m$.
- $C_{ik} C_{jk} D_{ij} x_k - x_k D_{ij} + C_{jk} x_j D_{ik} - C_{ij} D_{ik} x_j + D_{jk} x_i - C_{ij} C_{ik} x_i D_{jk} = 0$, $\forall 1 \leq i < j < k \leq m$ (non-degeneracy conditions).

If A is a G-algebra, then the set $\{x_j x_i - C_{ij} x_i x_j - D_{ij}, 1 \leq i < j \leq m\}$ is a Gröbner basis for the ideal it generates in $\mathbb{K}\langle x_1, \dots, x_m \rangle$ [16]. Also, if A is an algebra with PBW basis, then the non-degeneracy conditions are automatically satisfied.

Let \mathcal{B}' be the subalgebra of $U_{r,s}(\mathfrak{sl}_3)$ generated by $\{f_1, f_2, \omega'_1, \omega'_2\}$. Adding the element $\mathcal{F}_{21} = f_2 f_1 - s f_1 f_2$ to the generating set, \mathcal{B}' is generated by $\{f_1, \mathcal{F}_{21}, f_2, \omega'_1, \omega'_2\}$ subject to the relations

1. $\mathcal{F}_{21}f_1 = rf_1\mathcal{F}_{21}$ and $f_2\mathcal{F}_{21} = r\mathcal{F}_{21}f_2$,
2. $f_2f_1 = sf_1f_2 + \mathcal{F}_{21}$,
3. $\omega'_1\mathcal{F}_{21} = s^{-1}\mathcal{F}_{21}\omega'_1$ and $\omega'_2\mathcal{F}_{21} = r\mathcal{F}_{21}\omega'_2$,
4. the second type of relations (R3) from Definition II.1,
 - (a) $\omega'_1f_1 = rs^{-1}f_1\omega'_1$,
 - (b) $\omega'_2f_1 = sf_1\omega'_2$,
 - (c) $\omega'_1f_2 = r^{-1}f_2\omega'_1$,
 - (d) $\omega'_2f_2 = rs^{-1}f_2\omega'_2$, and
5. $\omega'_1\omega'_2 = \omega'_2\omega'_1$.

Therefore \mathcal{B}' is generated by $\{x_1 = f_1, x_2 = \mathcal{F}_{21}, x_3 = f_2, x_4 = \omega'_1, x_5 = \omega'_2\}$, subject to relations $\{x_jx_i = C_{ij}x_ix_j + D_{ij}, 1 \leq i < j \leq 5\}$ where the coefficients C_{ij} and polynomials D_{ij} are given by the relations above; that is $D_{ij} = 0$ if $(i, j) \neq (1, 3)$ and

1. $C_{12} = r$ and $C_{23} = r$,
2. $C_{13} = s$ and $D_{13} = \mathcal{F}_{21}$,
3. $C_{24} = s^{-1}$ and $C_{25} = r$,
4. (a) $C_{14} = rs^{-1}$,
- (b) $C_{15} = s$,
- (c) $C_{34} = r^{-1}$,
- (d) $C_{35} = rs^{-1}$, and
5. $C_{45} = 1$.

Recall from Chapter II that $\{f_1^{\alpha_1} \mathcal{F}_{21}^{\alpha_2} f_2^{\alpha_3} (\omega'_1)^{\alpha_4} (\omega'_2)^{\alpha_5}\}$ is a PBW basis for \mathcal{B}' ; hence the non-degeneracy conditions are satisfied. If we take $<_{\mathcal{B}'}$ to be the degree lexicographic order with $f_1 > \mathcal{F}_{21} > f_2 > \omega'_1 > \omega'_2$, then $\text{lm}(D_{13}) = \mathcal{F}_{21} < f_1 f_2$ since $\deg(\mathcal{F}_{21}) = 1 < 2 = \deg(f_1 f_2)$. Hence \mathcal{B}' is a G -algebra. Let I be the two-sided ideal of \mathcal{B}' generated by the set

$$\{(\omega'_1)^\ell - 1, (\omega'_2)^\ell - 1, f_1^\ell, \mathcal{F}_{21}^\ell, f_2^\ell\},$$

we have that $H = (\mathfrak{b}')^{\text{coop}} = \mathcal{B}'/I$.

2. The code

I now present how I defined \mathfrak{b}' in SINGULAR::PLURAL. The input and output are displayed in typewriter font and the output begins with the SINGULAR comment characters (`//`). For simplicity I wrote `W(i)` for ω'_i and `Q` for θ . The library `linalg.lib` contains the function `mat_rk` that calculates the rank of a matrix; from the library `matrix.lib` I use the command `gauss_col` which transforms a matrix into its column-reduced Gauss normal form. The library `qhmoduli.lib` contains the functions `Max` and `Min` which compute the maximum and minimum of a list of integers.

```
LIB "linalg.lib";
LIB "matrix.lib";
LIB "qhmoduli.lib";
```

For ℓ , y and z positive integers with $\gcd(y - z, \ell) = 1$, I define the ring B . I write the code in terms of parameters `l`, `y` and `z`; the values of these parameters can be fixed in a preamble as will be shown in Example III.3.

```
ring B = (0,Q), (F(1), F(21), F(2), W(1), W(2)), Dp;
minpoly = rootofUnity(1);
```

The underlying coefficient field has characteristic 0 and it contains \mathbb{Q} , which is a primitive ℓ th root of unity and is generated by the elements $F(1)$, $F(21)$, $F(2)$, $L(1)$, $L(2)$ (which correspond to $f_1, \mathcal{F}_{21}, f_2, \omega'_1$ and ω'_2 respectively). The monomial ordering Dp is the degree lexicographical order. I write the elements C_{ij} and D_{ij} that define the relations in \mathcal{B}' ; these are given with upper-triangular matrices C and D , and only the non-zero elements need to be given.

```
matrix C[5][5];
matrix D[5][5];
C[1,2] = Q^y; C[1,3] = Q^z; C[1,4] = Q^(y-z); C[1,5] = Q^z;
C[2,3] = Q^y; C[2,4] = Q^(-z); C[2,5] = Q^y;
C[3,4] = Q^(-y); C[3,5] = Q^(y-z);
C[4,5] = 1;
D[1,3] = F(21);
```

The command `ncalgebra(C,D)` creates the G -algebra with the relations given by C and D , and sets it as the base ring. I then give the generators of the ideal I .

```
ncalgebra(C,D);
option(redSB); option(redTail);
ideal I = F(1)^1, F(2)^1, W(1)^1 - 1, W(2)^1 - 1, (F(21))^1;
qring B = twostd(I);
```

The last command sets the base ring to be the quotient of the previous ring by the ideal I (the ideal has to be given by a two-sided Gröbner basis, and so I applied `twostd` to it). We now have \mathfrak{b}' as the base ring. The option `redSB` forces SINGULAR to work with reduced Gröbner basis, and `redTail` forces the reduction of the tails of polynomials during Gröbner basis computations. Next I describe how I generate the simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_3)$ -modules. Combining the definition of the \bullet_β action (Equation (I.3) in

Lemma I.29), together with the coproduct formulas in $H = (\mathbf{b}')^{\text{coop}}$ we have that for all $x \in H$ and $g \in G(H)$,

$$f_i \bullet_\beta x = -x S^{\text{op}}(f_i) + \beta(\omega'_i) f_i x (\omega'_i)^{-1} = -x f_i (\omega'_i)^{-1} + \beta(\omega'_i) f_i x (\omega'_i)^{-1} \quad (\text{III.1})$$

and

$$\omega'_i \bullet_\beta g = \beta(\omega'_i) \omega'_i g (\omega'_i)^{-1} = \beta(\omega'_i) g.$$

The second equation shows that if $g \in G(H)$, then $H \bullet_\beta g$ is generated by

$$\{(f_1^k \mathcal{F}_{21}^t f_2^m) \bullet_\beta g : 0 \leq k, t, m < \ell\}.$$

Recall from Chapter II that

$$\mathcal{F}_\ell = \{f_1^k \mathcal{F}_{21}^t f_2^m : 0 \leq k, t, m < \ell\}$$

and so

$$H \bullet_\beta g = \mathbb{K}\{f \bullet_\beta g : f \in \mathcal{F}_\ell\}.$$

Using Equation (III.1) I define the procedures `Beta1` and `Beta2`, so that `Beta1(a, h)` gives $f_1 \bullet_\beta h$ if $\beta(f_1) = \theta^a$ and `Beta(b, h)` gives $f_2 \bullet_\beta h$ if $\beta(f_2) = \theta^b$. Since $\mathcal{F}_{21} = f_2 f_1 - s f_1 f_2$, I define the procedure `Beta21` from the previous ones. For the results to be linear combinations of monomials where each generator appears as a factor at most ℓ times, I have to reduce the answer with respect to the ideal `std(0)`.

```

proc Beta1(int a, poly h)
{poly X;
  X = reduce((-h)*F(1)*W(1)^(1-1) + Q^a*F(1)*h*W(1)^(1-1), std(0));
  return(X);}

proc Beta2(int b, poly h)

```

```
{poly X;
  X = reduce((-h)*F(2)*W(2)^(1-1) + Q^b*F(2)*h*W(2)^(1-1), std(0));
  return(X);}
```

```
proc Beta21(int a, int b, poly h)
{return(Beta2(b, Beta1(a,h)) - Q^(z) * Beta1(a,Beta2(b,h)));}
```

Using compositions of these last procedures, I define the procedures PBeta1, PBeta2 and PBeta21, so that if $k \in \mathbb{N}$, $h \in H$ and $\beta(f_1) = \theta^a$ then PBeta1(a,h, k) gives $f_1^k \bullet_\beta h$, and similarly for $f_2^k \bullet_\beta h$ and $\mathcal{F}_{21}^k \bullet_\beta h$.

```
proc PBeta1(int a, poly h, int k)
{ poly Y = h;
  for(int n=1;n<=k;n++)
  { Y = Beta1( a, Y);}
  return(Y); }
```

```
proc PBeta2(int b, poly h, int k)
{ poly Y = h;
  for(int n=1;n<=k;n++)
  { Y = Beta2( b, Y);}
  return(Y);}
```

```
proc PBeta21(int a, int b, poly h, int k)
{ poly Y = h;
  for(int n=1;n<=k;n++)
  { Y = Beta21( a, b, Y);}
```

```
return(Y);}
```

Combining these procedures I define the procedure **Beta** so that if $0 \leq k, t, m < \ell$, $h \in H$ and $\beta : H \rightarrow \mathbb{K}$ is an algebra map given by $\beta(f_1) = \theta^a$ and $\beta(f_2) = \theta^b$, then $\text{Beta}(a, b, k, t, m, h)$ gives $(f_1^k \mathcal{F}_{21}^t f_2^m) \bullet_\beta h$.

```
proc Beta( int a , int b , int k, int t, int m, poly h)
{return( PBeta1( a, PBeta21( a, b, PBeta2(b,g,m) , t), k)) ;}
```

Fix a group-like element $g = (\omega'_1)^c (\omega'_2)^d \in H$. In what follows I will construct a basis and compute the dimensions for the module $H \bullet_\beta g$, where $\beta(\omega'_1) = \theta^{(z-y)c+yd}$ and $\beta(\omega'_2) = \theta^{-zc+(z-y)d}$. The basic idea is to consider the linear map $T_\beta : \mathbb{K}\mathcal{F}_\ell \rightarrow H$ given by $T_\beta(f) = f \bullet_\beta g$, and construct the matrix M representing T_β in the basis \mathcal{F}_ℓ and $\{fh : f \in \mathcal{F}_\ell, h \in G(H)\}$ of $\mathbb{K}\mathcal{F}_\ell$ and H respectively. Then $\dim(H \bullet_\beta g) = \text{rank}(M)$, and the non-zero columns of the column-reduced Gauss normal form of M give the coefficients for the elements of a basis of $H \bullet_\beta g$. The problem with this method is that since $\dim(H) = \ell^5$ and $\dim(\mathbb{K}\mathcal{F}_\ell) = \ell^3$, the size of M is $\ell^5 \times \ell^3$. Computing the Gauss normal form of these matrices is an expensive calculation even for small values of ℓ such as $\ell = 5$. However, by some reordering of \mathcal{F}_ℓ and of the PBW basis of H , M is block diagonal. I proceed to show how this is done.

For a monomial $h = f_1^{\alpha_1} \mathcal{F}_{21}^{\alpha_2} f_2^{\alpha_3} (\omega'_1)^{\alpha_5} (\omega'_2)^{\alpha_6}$ let $\deg_1(h) = \alpha_1 + \alpha_2$ and $\deg_2(h) = \alpha_2 + \alpha_3$. Note that Equation (III.1) implies that $h \bullet_\beta x$ is a linear combination of monomials m with $\deg_i(m) = \deg_i(h) + \deg_i(x)$. For all $0 \leq u, v < 2\ell$, let

$$D_{(u,v)} = \{h \in \mathcal{F}_\ell : \deg_1(h) = u \text{ and } \deg_2(h) = v\}$$

and

$$R_{(u,v)} = \{f(\omega'_1)^{-u} (\omega'_2)^{-v} g : f \in D_{(u,v)}\}.$$

Then for all $h \in D_{(u,v)}$, $h \bullet_{\beta} g \in \mathbb{K}R_{(u,v)}$. The possible pairs (u, v) are such that $0 \leq u, v \leq 2(\ell - 1)$ and since $|v - u|$ is the maximum power of \mathcal{F}_{21} that can be a factor of a monomial in $D_{(u,v)}$, we must have $|v - u| \leq \ell - 1$; that is $u - (\ell - 1) \leq v \leq u + \ell - 1$. Another way of describing the sets $D_{(u,v)}$ and $R_{(u,v)}$ is as follows.

$$\begin{aligned} D_{(u,v)} &= \{f_1^{u-i} \mathcal{F}_{21}^i f_2^{v-i}, \forall i \in \mathbb{N} : 0 \leq u - i, i, v - i \leq \ell - 1\} \\ &= \{f_1^{u-i} \mathcal{F}_{21}^i f_2^{v-i}, \forall i \in \mathbb{N} : n_{u,v} \leq i \leq m_{u,v}\} \end{aligned}$$

where $n_{u,v} = \max(0, \ell - 1 - u, \ell - 1 - v)$ and $m_{u,v} = \min(\ell - 1, u, v)$. Since $(\omega'_i)^{-1} = (\omega'_i)^{\ell-1}$, if $g = (\omega'_1)^c (\omega'_2)^d$ we also have

$$R_{(u,v)} = \{f(\omega'_1)^{(\ell-1)u+c} (\omega'_2)^{(\ell-1)v+d} : f \in D_{(u,v)}\}.$$

Remark III.2. It is clear that $\mathcal{F}_{\ell} = \bigcup D_{(u,v)}$, the union disjoint, and that $H \bullet_{\beta} g = \bigoplus \mathbb{K}R_{(u,v)}$. Therefore a basis for $H \bullet_{\beta} g$ is a disjoint union of the bases for $\mathbb{K}D_{(u,v)} \bullet_{\beta} g$ for all possible pairs (u, v) , and $\dim(H \bullet_{\beta} g) = \sum_{(u,v)} \dim(\mathbb{K}D_{(u,v)} \bullet_{\beta} g)$.

For ideals I_1 and I_2 given by a list of their generators, the command `coeffs` applied to the pair (I_1, I_2) returns a matrix A such that $I_2 A = I_1$, where the ideals I_1 and I_2 are thought of as one-row matrices whose entries are their generators. Therefore, for given u and v , if $M_{u,v}$ is the result of applying `coeffs` to the pair $(D_{(u,v)}, R_{(u,v)})$, then $\text{rank}(M_{(u,v)}) = \dim(\mathbb{K}D_{(u,v)} \bullet_{\beta} g)$, and if $N_{(u,v)}$ is the column-reduced Gauss normal form of $M_{(u,v)}$, the non-zero columns of $D_{(u,v)} N_{(u,v)}$ form a basis of $\mathbb{K}D_{(u,v)} \bullet_{\beta} g$.

I define the procedure `Submod`, where the output of `Submod(a, b, u, v)` is a list L , where the first component of the list is a basis for $D_{(u,v)} \bullet_{\beta} g$ and the second component is $\dim(D_{(u,v)} \bullet_{\beta} g)$.

```
proc Submod(int c, int d, int u, int v)
{ list L;
```



```

ideal D;
ideal R;
list e = u-(l-1),v-(l-1),0; int n= Max(e);
list f = u,v, l-1; int m= Min(f);
int a = (z-y)*c+ y*d; int b= -z*c+(z-y)*d;
for(int i= n; i<= m; i++)
{
D[i+1-n] = Beta(a, b , u-i, i, v-i , W(1)^c * W(2)^d);
R[i+1-n] = F(1)^(u-i)* F(2)^i* F(2)^(v-i)*
          W(1)^(((l-1)*u+c) mod l)* W(2)^(((l-1)*v+d) mod l);}
matrix M = coeffs(D,R);
matrix N = gauss_col(M);
matrix K[1][m-n+1] = R;
matrix S = K*N;
L[1] = compress(S);
L[2] = mat_rk(N);
return(L);}

```

The command `compress` deletes the zero columns of a matrix. For $g = (\omega'_1)^c(\omega'_2)^d$ the procedure `Totalbasis(c,d)` returns $\dim(H_{\bullet\beta}g)$ and a basis for $H_{\bullet\beta}g$, and is defined using Remark III.2.

```

proc Totalbasis(int c , int d)
{ list T; matrix A; int t; t = 0;
  for(int u = 0; u<=2*(l-1); u++)
    { list e = 0, u-(l-1);
      list f = u+(l-1), 2*(l-1);

```

```

    for(int v = Max(e); v <= Min(f); v++)
        { list M = Submod(c,d, u,v);
          A = compress(concat(A, M[1]));
          t = t + M[2];
        }
    }
T[1] = A; T[2] = t; return(T);
}

```

Example III.3. For $\ell = 5$, $y = 1$ and $z = 4$, for $g = (\omega'_1)^4(\omega'_2)^2$, I construct the module $H \bullet_{\beta} g$ as follows. To give SINGULAR:PLURAL the values of ℓ , y and z , I write at the beginning of the code

```

ring r0 = 0,x,dp;
int l = 1;
int y = 4;
int z = 1;

```

Then the command

```
Totalbasis(4,2);
```

returns

```

// [1]:
//   _[1,1]=W(1)^4*W(2)^2
//   _[1,2]=F(1)*W(1)^3*W(2)^2
//   _[1,3]=(-Q^3-Q^2-2*Q-1)*F(1)*F(2)*W(1)^3*W(2)+F(21)*W(1)^3*W(2)
// [2]:
//     3

```

which tells us that $\dim(H_{\bullet\beta}((\omega'_1)^4(\omega'_2)^2)) = 3$. In this case $\beta(\omega'_i) = \theta^{3.4+2} = \theta^4$ and $\beta(\omega'_2) = \theta^{-4.4+3.2} = 1$. A basis for $H_{\bullet\beta}g$ is $\{1_{\bullet\beta}g, f_{1\bullet\beta}g, \mathcal{F}_{21\bullet\beta}g\}$ since

$$\text{Beta}(4,0,0,0,0,W(1)^4*W(2)^2);$$

$$\text{Beta}(4,0,1,0,0,W(1)^4*W(2)^2)/(-Q^3-Q^2-2*Q-1);$$

$$\text{Beta}(4,0,0,1,0,W(1)^4*W(2)^2)/(-Q^3-Q^2-2*Q-1);$$

returns

$$// W(1)^4*W(2)^2$$

$$// F(1)*W(1)^3*W(2)^2$$

$$// (-Q^3-Q^2-2*Q-1)*F(1)*F(2)*W(1)^3*W(2)+F(21)*W(1)^3*W(2)$$

3. Computational results and conjectures

For $\ell = 5$, y and z such that $\gcd(3(y^2 - yz + z^2)(y - z), \ell) = 1$ and $g = (\omega'_1)^c(\omega'_2)^d$ ($0 \leq c, d < 5$) the corresponding $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_3)}$ -module $H_{\bullet\beta}g$ has dimension $\dim(c, d)$, where $\dim(c, d)$ is the entry in position $(c + 1, d + 1)$ of the symmetric matrix:

$$\text{DIM} = \begin{pmatrix} 1 & 60 & 90 & 15 & 18 \\ 60 & 8 & 10 & 15 & 39 \\ 90 & 10 & 19 & 35 & 3 \\ 15 & 15 & 35 & 63 & 6 \\ 18 & 39 & 3 & 6 & 125 \end{pmatrix}.$$

For $\ell = 7$, the results are analogous to the case $\ell = 5$, with matrix

$$\text{DIM} = \begin{pmatrix} 1 & 105 & 162 & 210 & 24 & 42 & 33 \\ 105 & 8 & 10 & 273 & 21 & 36 & 75 \\ 162 & 10 & 27 & 35 & 28 & 63 & 114 \\ 210 & 273 & 35 & 37 & 71 & 3 & 6 \\ 24 & 21 & 28 & 71 & 117 & 154 & 15 \\ 42 & 36 & 63 & 3 & 154 & 215 & 15 \\ 33 & 75 & 114 & 6 & 15 & 15 & 343 \end{pmatrix}.$$

By looking at these results, and the results obtained for other values of ℓ , I formulate the following conjecture:

Conjecture III.4. *Let y and z be integers such that $\gcd(3(y^2 - yz + z^2)(y - z), \ell) = 1$ and set $r = \theta^y$ and $s = \theta^z$. For integers $0 \leq c, d < \ell$ let $g = (\omega'_1)^c(\omega'_2)^d \in G(H)$ and $\beta : H \rightarrow \mathbb{K}$ be the algebra map given by $\beta(f_1) = \theta^{(z-y)c+d}$ and $\beta(f_2) = \theta^{-zc+(z-y)d}$. Let m_1 and m_2 be defined by*

$$m_1 \equiv (2c - d + 1) \text{ mod } \ell, \quad m_2 \equiv (2d - c + 1) \text{ mod } \ell \quad \text{and} \quad 0 < m_i \leq \ell.$$

If $m_1 + m_2 \leq \ell$ then

$$\dim(H_{\bullet} \beta g) = \frac{m_1 m_2 (m_1 + m_2)}{2}.$$

If $m_1 + m_2 > \ell$, let $x = m_1 + m_2 - \ell$, then

$$\dim(H_{\bullet} \beta g) = \frac{m_1 m_2 (m_1 + m_2)}{2} - \frac{(m_1 - x)(m_2 - x)(m_1 + m_2 - 2x)}{2}.$$

In the particular case when $y = 1$ and $z = \ell - 1$, the formulas above for the dimensions of the simple $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_3)$ -modules appeared in a work by Dobrev [8], where he calculated the dimensions of the simple modules for $U_{\theta}(\mathfrak{sl}_3)$, the infinite dimen-

sional one-parameter quantum group. By analyzing the results of the calculations in SINGULAR::PLURAL I formulate the following conjecture about simple $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_3)$ -modules.

Conjecture III.5. *For $g = (\omega'_1)^c(\omega'_2)^d \in G(H)$, let $m_1 \equiv (2c - d + 1) \bmod \ell$ and $m_2 \equiv (2d - c + 1) \bmod \ell$, $0 < m_i \leq \ell$. Let $\beta : H \rightarrow \mathbb{K}$ be the algebra map defined by $\beta(f_1) = \theta^{-2c+d} = \theta^{-m_1+1}$ and $\beta(f_2) = \theta^{-c+2d} = \theta^{-m_2+1}$ so that $H_{\bullet\beta}g$ is a $\overline{\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_3)}$ -module.*

If $m_1 + m_2 \leq \ell$, then the set

$$\{f_1^i \mathcal{F}_{21}^j f_2^k \bullet_{\beta} g : 0 \leq i < m_1, 0 \leq j < \ell, 0 \leq k < m_2 \text{ and } i + j + k \leq m_1 + m_2 - 2\}$$

is a basis for $H_{\bullet\beta}g$.

The conjecture was checked in PLURAL for $\ell = 5, 7, 11$, and calculations show that it holds when $m_2 = 1$.

CHAPTER IV

POINTED HOPF ALGEBRAS OF RANK ONE

Recently Andruskiewitsch and Schneider classified the pointed Hopf algebras with abelian groups of group-like elements, over an algebraically closed field of characteristic 0 [1]. Earlier, in 2005, Krop and Radford classified the pointed Hopf algebras of rank one, where $\text{rank}(H)+1$ is the rank of $H_{(1)}$ as an $H_{(0)}$ -module and H is generated by $H_{(1)}$ as an algebra, where $H_{(1)}$ is the first term of the coradical filtration of H [15]. They also studied the representation theory of $D(H)$ in a fundamental case. Using Radford's construction of simple modules, in Theorem IV.18, I give necessary and sufficient conditions for the tensor product of two $D(H)$ -modules to be completely reducible.

1. Pointed Hopf algebras of rank one of nilpotent type

Let G be a finite abelian group, \mathbb{K} an algebraically closed field of characteristic zero, $\chi : G \rightarrow \mathbb{K}$ a character and $a \in G$; we call the triple $\mathcal{D} = (G, \chi, a)$ *data*. Let $\ell := |\chi(a)|$, $N := |a|$ and $M = |\chi|$; note that ℓ divides both N and M . In [15] Krop and Radford defined the following Hopf algebra.

Definition IV.1. Let $\mathcal{D} = (G, \chi, a)$ be data. The Hopf algebra $H_{\mathcal{D}}$ is generated by G and x as a \mathbb{K} -algebra, with relations:

1. $x^{\ell} = 0$.
2. $xg = \chi(g)gx$, for all $g \in G$.

The coalgebra structure is given by $\Delta(x) = x \otimes a + 1 \otimes x$ and $\Delta(g) = g \otimes g$ for all $g \in G$.

The Hopf algebra $H_{\mathcal{D}}$ is pointed of rank one. Let $\Gamma = \text{Hom}(G, \mathbb{K}^{\times})$, the set of group homomorphisms from G to \mathbb{K}^{\times} also written \widehat{G} .

Proposition IV.2 (Krop and Radford [15]). *As a \mathbb{K} -algebra, $H_{\mathcal{D}}^*$ is generated by Γ and ξ subject to relations:*

1. $\xi^{\ell} = 0$.
2. $\xi\gamma = \gamma(a)\gamma\xi$, for all $\gamma \in \Gamma$.

The coalgebra structure of $H_{\mathcal{D}}^$ is determined by $\Delta(\xi) = \xi \otimes \chi + 1 \otimes \xi$ and $\Delta(\gamma) = \gamma \otimes \gamma$ for all $\gamma \in \Gamma$.*

Proposition IV.3 (Krop and Radford [15]). *The double $D(H_{\mathcal{D}})$ is generated by G , x , Γ , ξ subject to the relations defining $H_{\mathcal{D}}$ and $H_{\mathcal{D}}^*$ and the following relations:*

1. $g\gamma = \gamma g$ for all $g \in G$ and $\gamma \in \Gamma$.
2. $\xi g = \chi^{-1}(g)g\xi$ for all $g \in G$.
3. $[x, \xi] = a - \chi$.
4. $\gamma(a)x\gamma = \gamma x$ for all $\gamma \in \Gamma$.

Recall that the coalgebra structure of $H_{\mathcal{D}}^*$ in $D(H_{\mathcal{D}})$ is the co-opposite to the one in H^* . Then in $D(H_{\mathcal{D}})$, $\Delta(\xi) = \chi \otimes \xi + \xi \otimes 1$. Note that $H_{\mathcal{D}}$ satisfies the hypothesis of Proposition I.30, where elements in G have degree 0 and x has degree 1. Therefore, simple $D(H_{\mathcal{D}})$ -modules are of the form $H \bullet_{\beta} g$, for $g \in G$ and $\beta \in G(H^*) = \Gamma$.

2. Factorization of simple $D(H_{\mathcal{D}})$ -modules

In this section I study under what conditions a simple $D(H_{\mathcal{D}})$ -module can be factored as the tensor product of a one-dimensional module with a simple module which is also a module for $\overline{D(H_{\mathcal{D}})} = D(H_{\mathcal{D}})/D(H_{\mathcal{D}})(\mathbb{K}G_C(D(H_{\mathcal{D}})))^+$. I also study, under certain

conditions on the parameters, the reducibility of the tensor product of two simple $D(H_{\mathcal{D}})$ -modules.

I start by describing the central group-like elements of $D(H_{\mathcal{D}})$. It is clear that $G(D(H_{\mathcal{D}})) = G \times \Gamma$. An element $(g, \gamma) \in G \times \Gamma$ will be denoted by $g\gamma$. An element $g\gamma$ is central in $D(H_{\mathcal{D}})$ if and only if $(g\gamma)x = x(g\gamma)$ and $(g\gamma)\xi = \xi(g\gamma)$. Using the relations of $D(H_{\mathcal{D}})$, we have that

$$g\gamma x = \gamma(a)gx\gamma = \chi^{-1}(g)\gamma(a)xg\gamma,$$

and

$$g\gamma\xi = \gamma g\xi = \chi(g)\gamma\xi g = \chi(g)\gamma(a)^{-1}\xi\gamma g.$$

Hence, $g\gamma$ is central if only if $\chi^{-1}(g)\gamma(a) = 1$. Let $\text{ev}_{\chi^{-1}a} : G \times \Gamma \rightarrow \mathbb{K}^{\times}$ be the character given by $\text{ev}_{\chi^{-1}a}(g\gamma) = \chi^{-1}(g)\gamma(a)$; we just showed the following lemma:

Lemma IV.4. $G_C(D(H_{\mathcal{D}})) = \text{Ker}(\text{ev}_{\chi^{-1}a})$.

For $\alpha : D(H_{\mathcal{D}}) \rightarrow \mathbb{K}$ an algebra map, let \mathbb{K}_{α} be the one-dimensional module defined by $h \cdot k = \alpha(h)k$ for all $h \in D(H_{\mathcal{D}})$ and $k \in \mathbb{K}$. Note that α being an algebra map implies that $\alpha(x) = \alpha(\eta) = 0$ (because $0 = x^{\ell} = \xi^{\ell}$) and $\alpha(a) = \alpha(\chi)$ (by the third relation in Proposition IV.3). Since $\alpha(x) = \alpha(\eta) = 0$, we can think of α as a group homomorphism $\alpha : G \times \Gamma \rightarrow \mathbb{K}^{\times}$, that is, $\alpha \in \widehat{G \times \Gamma} \simeq \Gamma \times G$. Let $\beta_{\alpha} \in \Gamma$ and $g_{\alpha} \in G$ so that $\alpha = \beta_{\alpha}g_{\alpha}$; that is $\alpha(g\gamma) = \beta_{\alpha}(g)\gamma(g_{\alpha})$ for all $g\gamma$ in $G \times \Gamma$. If we extend β_{α} to $H_{\mathcal{D}}$ by setting $\beta_{\alpha}(x) = 0$ and also call this extension β_{α} (as no confusion will arise), we have $\beta_{\alpha} = \alpha|_{H_{\mathcal{D}}}$.

Proposition IV.5. $\mathbb{K}_{\alpha} \simeq H_{\mathcal{D}\bullet\beta_{\alpha}g_{\alpha}}$ as Yetter-Drinfel'd $H_{\mathcal{D}}$ -modules.

Proof. Since \mathbb{K}_{α} is a simple Yetter-Drinfel'd module, there exists an isomorphism of Yetter-Drinfel'd modules $\Phi : \mathbb{K}_{\alpha} \rightarrow H_{\mathcal{D}\bullet\beta g}$ for some algebra map $\beta : H_{\mathcal{D}} \rightarrow \mathbb{K}$ and

some $g \in G$. We may assume that $\Phi(1) = g$. Let $h \in G$, we have

$$h \bullet_{\beta} g = \beta(h)g.$$

Since Φ is a module map,

$$\begin{aligned} h \bullet_{\beta} g &= h \bullet_{\beta} \Phi(1) = \Phi(h \cdot 1) = \Phi(\alpha(h)) \\ &= \alpha(h)\Phi(1) = \beta_{\alpha}(h)g. \end{aligned}$$

We then have $\beta(h) = \beta_{\alpha}(h)$ for all h in G , and since $\beta(x) = \beta_{\alpha}(x) = 0$, $\beta = \beta_{\alpha}$.

If $\gamma \in \Gamma$, then

$$\gamma \bullet_{\beta} g = \gamma(g)g.$$

On the other hand,

$$\gamma \bullet_{\beta} g = \gamma \bullet_{\beta} \Phi(1) = \Phi(\gamma \cdot 1) = \Phi(\alpha(\gamma)1) = \alpha(\gamma)\Phi(1) = \gamma(g_{\alpha})g.$$

Then $\gamma(g) = \gamma(g_{\alpha})$ for all $\gamma \in \Gamma$, hence $g = g_{\alpha}$. □

For simplicity let $K = \text{Ker}(ev_{\chi^{-1}a})$. If $\alpha = \beta_{\alpha}g_{\alpha} \in \Gamma \times G$, the condition $\alpha(a) = \alpha(\chi)$ is $\beta_{\alpha}(a) = \chi(g_{\alpha})$ or $\chi^{-1}(g_{\alpha})\beta_{\alpha}(a) = 1$. Hence, α in $\Gamma \times G$ defines a one-dimensional module if and only if $g_{\alpha}\beta_{\alpha} \in \text{Ker}(ev_{\chi^{-1}a}) = K$. This, together with the previous proposition, shows

Corollary IV.6. *The set $\mathcal{S}_{D(H_{\mathcal{D}})}^1$ of isomorphism classes of one dimensional $D(H_{\mathcal{D}})$ -modules is in one to one correspondence with K .*

Recall that $\overline{D(H_{\mathcal{D}})} = D(H_{\mathcal{D}})/D(H_{\mathcal{D}})(\mathbb{K}G_C(D(H_{\mathcal{D}})))^+$. Since $G_C(D(H_{\mathcal{D}})) = K$,

$$D(H_{\mathcal{D}})(\mathbb{K}G_C(D(H_{\mathcal{D}})))^+ = D(H)\{g\gamma - 1 : g\gamma \in K\}.$$

For a group A and a subgroup $B \subset A$, let

$$B^\perp = \{f \in \widehat{A} : f(b) = 1 \text{ for all } b \in B\}.$$

Note that $K^\perp \subset \widehat{G \times \Gamma} \simeq \Gamma \times G$.

Proposition IV.7. *For $\beta \in G(H_{\mathcal{D}}^*) = \Gamma$ and $g \in G$, the simple $D(H_{\mathcal{D}})$ -module $H_{\mathcal{D}\bullet\beta}g$ is also a $\overline{D(H_{\mathcal{D}})}$ -module via the quotient map, if and only if $\beta g \in K^\perp$.*

Proof. $H_{\mathcal{D}\bullet\beta}g$ is a $\overline{D(H_{\mathcal{D}})}$ -module, if and only if $f\gamma \cdot (h\bullet\beta g) = h\bullet\beta g$, for all $f\gamma \in K$ and $h \in H_{\mathcal{D}}$. Since $K \subset \mathcal{Z}(D(H_{\mathcal{D}}))$, if $f\gamma \in K$ then $f\gamma \cdot (h\bullet\beta g) = (f\gamma h) \cdot g = (hf\gamma) \cdot g = h\bullet\beta((f\gamma) \cdot g)$. Thus, $H_{\mathcal{D}\bullet\beta}g$ is a $\overline{D(H_{\mathcal{D}})}$ -module, if and only if $f\gamma \cdot g = g$, for all $f\gamma \in K$. Now $f\gamma \cdot g = f\bullet\beta\gamma(g)g = \gamma(g)\beta(f)g$. And so, $H_{\mathcal{D}\bullet\beta}g$ is a $\overline{D(H_{\mathcal{D}})}$ -module, if and only if $\gamma(g)\beta(f) = 1$ for all $f\gamma \in K$; that is, if and only if, $\beta g \in K^\perp$. \square

Lemma IV.8. $K^\perp = \langle \text{ev}_{\chi^{-1}a} \rangle$.

Proof. Since $K^\perp \simeq \widehat{\left(\frac{G \times \Gamma}{K}\right)}$, we have $|K^\perp| = \left|\frac{G \times \Gamma}{K}\right| = |\text{Im } \text{ev}_{\chi^{-1}a}| = |\text{ev}_{\chi^{-1}a}|$; the last equality holding as $\text{Im } \text{ev}_{\chi^{-1}a}$ is cyclic (since it is a finite subgroup of \mathbb{K}^\times). By the definitions of K and K^\perp , $\text{ev}_{\chi^{-1}a} \in K^\perp$, hence $K^\perp = \langle \text{ev}_{\chi^{-1}a} \rangle$. \square

It will be convenient to think of K^\perp as a subgroup of $G \times \Gamma$ via the identification $\widehat{G \times \Gamma} \simeq \widehat{G} \times \widehat{\Gamma} \simeq \Gamma \times G \simeq G \times \Gamma$. Under this identification we have $K^\perp = \langle a\chi^{-1} \rangle$.

Remark IV.9. We can restate Proposition IV.7 as follows: the simple $D(H_{\mathcal{D}})$ -modules that are also $\overline{D(H_{\mathcal{D}})}$ -modules are of the form $H_{\mathcal{D}\bullet(\chi^{-c})}a^c$, for $c = 1, \dots, |a\chi^{-1}|$.

Recall that $\mathcal{S}_{D(H_{\mathcal{D}})}$ denotes the set of isomorphism classes of simple $D(H_{\mathcal{D}})$ -modules. Combining Proposition I.33, Corollary IV.6 and Proposition IV.7, we get that the map

$$\Phi : \mathcal{S}_{\overline{D(H_{\mathcal{D}})}} \times \mathcal{S}_{D(H_{\mathcal{D}})}^1 \rightarrow \mathcal{S}_{D(H_{\mathcal{D}})}$$

given by $\Phi(U, V) = U \otimes V$, is equivalent to the multiplication map

$$\mu : K^\perp \times K \rightarrow G \times \Gamma,$$

under the identification of simple $D(H_{\mathcal{D}})$ -modules with elements of $G \times \Gamma$.

Theorem IV.10. *The map Φ as above is a bijection if and only if ℓ is odd and $\ell = M = N$.*

Proof. By the last remark, Φ is an bijection, if and only if $G \times \Gamma = K^\perp \times K$, that is $G \times \Gamma = K^\perp K$ and $K \cap K = \{1\}$. Now $|K^\perp| = \frac{|G \times \Gamma|}{|K|} = \frac{|G \times \Gamma|}{|K|}$, and so $|K^\perp K| = \frac{|K^\perp| |K|}{|K^\perp \cap K|} = \frac{|G \times \Gamma|}{|K^\perp \cap K|}$. We then have that $K^\perp K = G \times \Gamma$ if and only if $K^\perp \cap K = \{1\}$. If $\ell = M = N$, then $|a| = |\chi| = \ell$ and so $|a\chi^{-1}| = \ell$. Since $K^\perp \cap K \subset K^\perp = \langle a\chi^{-1} \rangle$, we have that $K^\perp \cap K = \langle (a\chi^{-1})^r \rangle$ for some $r \in \{1, \dots, \ell\}$. Since $(a\chi^{-1})^r \in K = \text{Ker}(\text{ev}_{\chi^{-1}a})$, $1 = \text{ev}_{\chi^{-1}a}((a\chi^{-1})^r) = (\chi^{-1}(a))^{2r}$ and so $\ell | 2r$. If ℓ is odd, then $\ell | r$ and so $(a\chi^{-1})^r = 1$, giving $K^\perp \cap K = \{1\}$.

Conversely, if $K^\perp \cap K = \{1\}$, let $n = |a\chi^{-1}|$. Then for all $r \in \{1, \dots, n-1\}$, $(a\chi^{-1})^r \notin K$. If either $M \neq \ell$ or $N \neq \ell$, then $n > \ell$ and so $(a\chi^{-1})^\ell \notin K$, which is a contradiction since $\text{ev}_{\chi^{-1}a}((a\chi^{-1})^\ell) = \chi^{-1}(a)^{2\ell} = 1$. Hence, $\ell = M = N$. If ℓ is even, then $(a\chi^{-1})^{\frac{\ell}{2}} \notin K$, which is again a contradiction since $\text{ev}_{\chi^{-1}a}((a\chi^{-1})^{\frac{\ell}{2}}) = \chi^{-1}(a)^\ell = 1$. Hence ℓ is odd. \square

Next I describe the structure of $\overline{D(H_{\mathcal{D}})}$ under the hypothesis of the last Theorem.

Proposition IV.11. *If ℓ is odd and $\ell = N = M$, then $\overline{D(H_{\mathcal{D}})} \simeq \mathfrak{u}_\theta(\mathfrak{sl}_2)$ as Hopf algebras, where $\theta = \chi(a)^{-\frac{1}{2}}$.*

Proof. Recall that $\mathfrak{u}_\theta(\mathfrak{sl}_2) = \mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2) / \langle (\omega'_1)^{-1} - \omega_1 \rangle$. Since there is only one generator of each kind, I will omit the subindex 1; we then have that $\mathfrak{u}_\theta(\mathfrak{sl}_2)$ is generated by e ,

f and ω , with relations:

$$e^\ell = 0 = f^\ell, \quad \omega^\ell = 1, \quad \omega e = \theta^2 e \omega, \quad \omega f = \theta^{-2} f \omega \quad \text{and} \quad [e, f] = \frac{1}{\theta - \theta^{-1}} \omega - \omega^{-1}.$$

In the proof of the previous proposition, we showed that if ℓ is odd and $\ell = N = M$, then $G \times \Gamma = \langle a\chi^{-1} \rangle K$, and so $\langle \chi^{-1} a \rangle$ is a complete set of representatives of the classes in $\frac{G \times \Gamma}{K}$. Let $\psi : D(H_D) \rightarrow \mathbf{u}_\theta(\mathfrak{sl}_2)$ be the algebra map such that

- $\psi(g\gamma) = \omega^{-2c}$ if $g\gamma \in (a\chi^{-1})^c K, \forall g\gamma \in G \times \Gamma$,
- $\psi(\xi) = e$ and
- $\psi(x) = (\theta - \theta^{-1})f$.

For ψ to be defined, it must commute with the defining relations of $D(H_D)$ (from Definition IV.1 and Propositions IV.2 and IV.3). This is the case by the following calculations:

1. $\psi(x)\psi(g) = \chi(g)\psi(g)\psi(x)$, for all $g \in G$:

Let $g \in G$; if $g \in (a\chi^{-1})^c K$, then $g = (a\chi^{-1})^c g_K \chi^c$, with $g_K \chi^c \in K$. Hence $\chi^c(a)\chi^{-1}(g_K) = 1$ and so $\chi(g_K) = \chi^c(a) = q^c$. Therefore

$$\chi(g) = \chi(a^c g_K) = \chi(a^c)\chi(g_K) = q^{2c}.$$

Then,

$$\begin{aligned} \psi(x)\psi(g) &= (\theta - \theta^{-1})f\omega^{-2c} = (\theta - \theta^{-1})\theta^{-4c}\omega^{-2c}f = \chi(g)\omega^{-2c}(\theta - \theta^{-1})f \\ &= \chi(g)\psi(g)\psi(x). \end{aligned}$$

2. $\psi(\xi)\psi(\gamma) = \gamma(a)\psi(\gamma)\psi(\xi)$, for all $\gamma \in \Gamma$:

Let $\gamma \in \Gamma$, in a similar way as in the previous relation, it can be shown that if

$\gamma \in (a\chi^{-1})^c K$, then $\gamma(a) = q^{-2c}$. We then have

$$\psi(\xi)\psi(\gamma) = e\omega^{-2c} = \theta^{4c}\omega^{-2c}e = \gamma(a)\psi(\gamma)\psi(\xi).$$

3. $[\psi(x), \psi(\xi)] = \psi(a) - \psi(\chi)$:

To prove this, we first need to know the images of a and χ under ψ . Since ℓ is odd, let $c \in \mathbb{Z}$ be such that $2c = 1 \pmod{\ell}$. Then, $a = (a\chi^{-1})^c(a\chi)^c$, and since $a\chi \in K$, we have that

$$\psi(a) = \omega^{-2c} = \omega^{-1}. \quad (\text{IV.1})$$

Similarly, $\chi = (a\chi^{-1})^{-c}(a\chi)^c$ and so $\psi(\chi) = \omega$. Now

$$\begin{aligned} [\psi(x), \psi(\xi)] &= (\theta - \theta^{-1})[f, e] = -(\theta - \theta^{-1})[e, f] = -\frac{\theta - \theta^{-1}}{\theta - \theta^{-1}}(\omega - \omega^{-1}) \\ &= \omega^{-1} - \omega = \psi(a) - \psi(\chi). \end{aligned}$$

Clearly $\psi(x)^\ell = 0 = \psi(\xi)^\ell$ and $\psi(g)\psi(\gamma) = \psi(\gamma)\psi(g)$ for all $g \in G$ and $\gamma \in \Gamma$. The other relations follow in a similar way as 1 and 2 above.

Next we need to show that ψ is a map of coalgebras. Group-like elements in $D(H_{\mathcal{D}})$ are mapped to group-like elements in $\mathfrak{u}_\theta(\mathfrak{sl}_2)$. Moreover,

$$\begin{aligned} \psi \otimes \psi(\Delta(x)) &= \psi \otimes \psi(x \otimes a + 1 \otimes x) = (\theta - \theta^{-1})(f \otimes \omega^{-1} + 1 \otimes f) \\ &= (\theta - \theta^{-1})\Delta(f) = \Delta(\psi(x)) \end{aligned}$$

and

$$\psi \otimes \psi(\Delta(\xi)) = \psi \otimes \psi(\chi \otimes \xi + \xi \otimes 1) = (\omega \otimes e + e \otimes 1) = \Delta(e) = \Delta(\psi(\xi)).$$

Therefore ψ is a map of Hopf algebras.

Recall that $D(H_{\mathcal{D}})(\mathbb{K}K)^+ = D(H_{\mathcal{D}})\{k - 1 : k \in K\}$. Note that $\psi(K) = \{1\}$ and so $\psi(\{k - 1 : k \in K\}) = 0$. Therefore $D(H_{\mathcal{D}})(\mathbb{K}K)^+ \subset \text{Ker}(\psi)$ and the map ψ

induces a Hopf algebra map $\overline{\psi} : \overline{D(H_{\mathcal{D}})} \rightarrow \mathfrak{u}_{\theta}(\mathfrak{sl}_2)$. Since ℓ is odd, $\langle \omega \rangle = \langle \omega^{-2} \rangle$, and so $\overline{\psi}$ is surjective.

By Remark I.15,

$$\begin{aligned} \dim(\overline{D(H_{\mathcal{D}})}) &= \frac{\dim(D(H_{\mathcal{D}}))}{\dim(\mathbb{K}K)} = \frac{|G \times \Gamma| \ell^2}{|K|} = |K^{\perp}| \ell^2 = |\langle a\chi^{-1} \rangle| \ell^2 = \ell^3 \\ &= \dim(\mathfrak{u}_{\theta}(\mathfrak{sl}_2)). \end{aligned}$$

Hence, $\overline{\psi}$ is an isomorphism. □

Remark IV.12. Let \mathfrak{b}' be (as in Chapter II) the subalgebra of $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2)$ generated by f and ω' and $H = (\mathfrak{b}')^{\text{coop}}$. Via the isomorphism $\overline{\psi}$ defined in the proof of Proposition IV.11, a simple $D(H_{\mathcal{D}})$ -module of the form $H_{\mathcal{D}^{\bullet}(\chi^{-c})}(a^c)$ is also a $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -module. Explicitly, for $h \in \mathfrak{u}_{\theta}(\mathfrak{sl}_2) = \overline{\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2)}$ and $m \in H_{\mathcal{D}^{\bullet}(\chi^{-c})}(a^c)$, $h \cdot m = \overline{\psi}^{-1}(h) \cdot m$. Therefore, as $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -modules, $H_{\mathcal{D}^{\bullet}(\chi^{-c})}(a^c) \simeq H_{\bullet\beta}(\omega'^d)$ with $\beta(\omega') = \theta^{-2d}$ for some $d \in \mathbb{Z}$. By analyzing the action of ω' on both of this modules, it follows that $d = -c$. Conversely, a simple $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -module $H_{\bullet\beta}(\omega')^d$ becomes a simple $D(H_{\mathcal{D}})$ -module via $\overline{\psi}$, and is isomorphic to $H_{\mathcal{D}^{\bullet}(\chi^d)}(a^{-d})$ as $D(H_{\mathcal{D}})$ -modules.

I finish this section by studying the reducibility of tensor products of simple $D(H_{\mathcal{D}})$ -modules when $n = M = N$ is odd.

In [19], Radford used his construction to describe simple modules for the Drinfel'd Double of the Taft algebra, which is isomorphic to $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2)$ when ℓ is odd (ℓ is the order of θ). Translating his result to our notation ($H = (\mathfrak{b}')^{\text{coop}}$, generated by ω' and f and the corresponding relations) we have

Proposition IV.13 (Radford [19]). *For $g = (\omega')^c$ and $\beta : H \rightarrow \mathbb{K}$ an algebra morphism, let $r \geq 0$ be minimal such that $\beta(\omega') = \theta^{2(c-r)}$. Then the simple $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2)$ -module $H_{\bullet\beta}g$ is $(r+1)$ -dimensional with basis $\{g, f_{\bullet\beta}g, \dots, f_c^r_{\bullet\beta}g\}$ and $f^{r+1}_{\bullet\beta}g = 0$.*

In [7], H-X. Chen studied the reducibility of tensor products of these simple modules:

Proposition IV.14 (Chen [7]). *Given $g = (\omega')^c$, $g' = (\omega')^{c'}$ in $G(H)$ and $\beta, \beta' \in G(H^*)$, let $r, r' \in \{0, \dots, \ell - 1\}$ be such that $\beta(\omega') = \theta^{2(c-r)}$ and $\beta'(\omega') = \theta^{2(c'-r')}$. Then the $\mathfrak{u}_{\theta, \theta^{-1}}(\mathfrak{sl}_2)$ -module $H_{\bullet\beta}g \otimes H_{\bullet\beta'}g'$ is completely reducible if and only if $r + r' < \ell$. Moreover, let*

$$g_j = gg'(\omega')^{-j} \quad \text{and} \quad \beta_j(\omega') = \theta^{2j}\beta(\omega')\beta'(\omega');$$

if $r + r' < \ell$ then

$$H_{\bullet\beta}g \otimes H_{\bullet\beta'}g' \simeq \bigoplus_{j=0}^{\min(r, r')} H_{\bullet\beta_j}g_j.$$

If $r + r' \geq \ell$, let $t = r + r' - \ell + 1$; then

$$\text{Soc}(H_{\bullet\beta}g \otimes H_{\bullet\beta'}g') \simeq \bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r, r')} H_{\bullet\beta_j}g_j.$$

Remark IV.15. By Example II.9, if $H_{\bullet\beta}(\omega')^c$ is naturally a $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -module, then $\beta = \beta_g$, i.e. $\beta(\omega') = \theta^{-2c} = \theta^{2(c-2c)}$. Then the number r from Proposition IV.13 is $r = 2c \bmod \ell$, with $0 \leq r < \ell$. I will denote such number by r_c .

We get the following corollary for simple $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -modules:

Corollary IV.16. *Given $g = (\omega')^c$ and $g' = (\omega')^{c'}$ in $G(H)$. If $r_c + r_{c'} < \ell$ then*

$$H_{\bullet\beta_g}g \otimes H_{\bullet\beta_{g'}}g' \simeq \bigoplus_{j=0}^{\min(r_c, r_{c'})} H_{\bullet\beta_j}g_j,$$

as $\mathfrak{u}_{\theta}(\mathfrak{sl}_2)$ -modules, where $g_j = gg'(\omega')^{-j}$ and $\beta_j = \beta_{g_j}$.

Remark IV.17. This last corollary is a particular case of a more general formula for simple modules for the non-restricted quantum group $U_q(\mathfrak{sl}_2)$, that appears as an exercise in [3].

We have an analogous result to Proposition IV.14 for $D(H_{\mathcal{D}})$ -modules:

Theorem IV.18. *If $\ell = M = N$ is odd and $g\beta, g'\beta' \in G \times \Gamma = G(D(H_{\mathcal{D}}))$, let c and $c' \in \mathbb{Z}$ such that $(a^{-1}\chi)^c$ and $(a^{-1}\chi)^{c'}$ are representatives of the classes of $g\beta$ and $g'\beta'$ in $G \times \Gamma/K$ respectively. Then the $D(H_{\mathcal{D}})$ -module $H_{\mathcal{D}^\bullet\beta}g \otimes H_{\mathcal{D}^\bullet\beta'}g'$ is completely reducible if and only if $r_c + r_{c'} < \ell$. Moreover, let*

$$g_j = gg'a^j \quad \text{and} \quad \beta = \chi^{-j}\beta\beta';$$

if $r_c + r_{c'} < \ell$ then

$$H_{\mathcal{D}^\bullet\beta}g \otimes H_{\mathcal{D}^\bullet\beta'}g' \simeq \bigoplus_{j=0}^{\min(r_c, r_{c'})} H_{\mathcal{D}^\bullet\beta_j}g_j.$$

If $r_c + r_{c'} \geq \ell$, then

$$\text{Soc}(H_{\mathcal{D}^\bullet\beta}g \otimes H_{\mathcal{D}^\bullet\beta'}g') \simeq \bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r_c, r_{c'})} H_{\mathcal{D}^\bullet\beta_j}g_j,$$

where $t = r_c + r_{c'} - \ell + 1$.

Proof. Let $g_K\beta_K$ and $g'_K\beta'_K \in K$ such that $g\beta = (a^{-1}\chi)^c g_K\beta_K$ and $g'\beta' = (a^{-1}\chi)^{c'} g'_K\beta'_K$. By Proposition IV.10, $H_{\mathcal{D}^\bullet\beta}g \simeq H_{\mathcal{D}^\bullet\chi^c}a^{-c} \otimes H_{\mathcal{D}^\bullet\beta_K}g_K$, the first factor in $\mathcal{S}_{\overline{D(H_{\mathcal{D}})}}$, and the second factor in $\mathcal{S}_{D(H_{\mathcal{D}})}^1$. Similarly $H_{\mathcal{D}^\bullet\beta'}g' = H_{\mathcal{D}^\bullet\chi^{c'}}a^{-c'} \otimes H_{\mathcal{D}^\bullet\beta'_K}g'_K$. Then

$$\begin{aligned} H_{\mathcal{D}^\bullet\beta}g \otimes H_{\mathcal{D}^\bullet\beta'}g' &\simeq (H_{\mathcal{D}^\bullet\chi^c}a^{-c} \otimes H_{\mathcal{D}^\bullet\beta_K}g_K) \otimes (H_{\mathcal{D}^\bullet\chi^{c'}}a^{-c'} \otimes H_{\mathcal{D}^\bullet\beta'_K}g'_K) \\ &\simeq (H_{\mathcal{D}^\bullet\chi^c}a^{-c} \otimes H_{\mathcal{D}^\bullet\chi^{c'}}a^{-c'}) \otimes (H_{\mathcal{D}^\bullet\beta_K}g_K \otimes H_{\mathcal{D}^\bullet\beta'_K}g'_K) \\ &\simeq (H_{\mathcal{D}^\bullet\chi^c}a^{-c} \otimes H_{\mathcal{D}^\bullet\chi^{c'}}a^{-c'}) \otimes H_{\mathcal{D}^\bullet_{\beta_K\beta'_K}}g_K g'_K; \end{aligned}$$

the second isomorphism by symmetry of tensor products of modules for $D(H_{\mathcal{D}})$, and the third by combining Propositions I.33 and I.34. Let $\gamma, \gamma' : H \rightarrow \mathbb{K}$ be the algebra maps given by $\gamma(\omega') = \theta^{-2c}$ and $\gamma'(\omega') = \theta^{-2c'}$. If $r_c + r_{c'} < \ell$, we have the following

isomorphisms of $\mathfrak{u}_\theta(\mathfrak{sl}_2)$ -modules:

$$H_{\mathcal{D}^\bullet \chi^c a^{-c}} \otimes H_{\mathcal{D}^\bullet \chi^{c'} a^{-c'}} \simeq H_{\bullet \gamma}(\omega')^c \otimes H_{\bullet \gamma'}(\omega')^{c'} \simeq \bigoplus_{j=0}^{\min(r, r')} H_{\bullet \beta_j} g_j,$$

where $g_j = (\omega')^{c+c'-j}$ and $\gamma_j(\omega') = \theta^{-2(c+c'-j)}$, the first isomorphism following from the Remark IV.12 and the second from Corollary IV.16. Again by the Remark IV.12, the j^{th} summand of the last module is isomorphic to $H_{\mathcal{D}^\bullet \chi^{-c_j} a^{c_j}}$ as $D(H_{\mathcal{D}})$ -modules, where $c_j = -(c + c' - j)$. Then

$$H_{\mathcal{D}^\bullet \beta} g \otimes H_{\mathcal{D}^\bullet \beta'} g' \simeq \left(\bigoplus_{j=0}^{\min(r, r')} H_{\mathcal{D}^\bullet \chi^{-c_j} a^{c_j}} \right) \otimes H_{\mathcal{D}^\bullet \beta_K \beta'_K} g_K g'_K \simeq \bigoplus_{j=0}^{\min(r, r')} H_{\mathcal{D}^\bullet \gamma_j} g_j,$$

where

$$g_j = a^{c_j} g_K g'_K = a^{-c} g_K a^{-c'} g'_K a^j = g g' a^j$$

and

$$\gamma_j = \chi^{-c_j} \beta_K \beta'_K = \chi^c \beta_K \chi^{c'} \beta'_K \chi^{-j} = \beta \beta' \chi^{-j}.$$

If $r_c + r_{c'} \geq \ell$, we have

$$H_{\mathcal{D}^\bullet \beta} g \otimes H_{\mathcal{D}^\bullet \beta'} g' \simeq \left(H_{\mathcal{D}^\bullet \chi^{-c} a^c} \otimes H_{\mathcal{D}^\bullet \chi^{-c'} a^{c'}} \right)_{\beta_K \beta'_K}$$

and by Remark I.34 we have

$$\text{Soc} \left(H_{\mathcal{D}^\bullet \beta} g \otimes H_{\mathcal{D}^\bullet \beta'} g' \right) \simeq \left(\text{Soc} \left(H_{\mathcal{D}^\bullet \chi^{-c} a^c} \otimes H_{\mathcal{D}^\bullet \chi^{-c'} a^{c'}} \right) \right)_{\beta_K \beta'_K}.$$

With a similar reasoning as before, we get that

$$\text{Soc} \left(H_{\mathcal{D}^\bullet \chi^{-c} a^c} \otimes H_{\mathcal{D}^\bullet \chi^{-c'} a^{c'}} \right) \simeq \bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r_c, r_{c'})} H_{\bullet \chi^{-c_j} a^{c_j}},$$

where $c_j = -(c + c' - j)$. Therefore

$$\begin{aligned}
\text{Soc}(H_{\mathcal{D}^\bullet\beta}g \otimes H_{\mathcal{D}^\bullet\beta'}g') &\simeq \left(\bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r_c, r_{c'})} H_{\chi^{-c_j}} a^{c_j} \right)_{\beta_K \beta'_K} \\
&\simeq \left(\bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r_c, r_{c'})} H_{\chi^{-c_j}} a^{c_j} \right) \otimes H_{\beta_K * \beta'_K} g_K g'_K \\
&\simeq \bigoplus_{j=\lceil \frac{t+1}{2} \rceil}^{\min(r_c, r_{c'})} H_{\beta_j} g_j,
\end{aligned}$$

where $g_j = h_j g_K g'_K = a^{-c-c'+j} g_K g'_K = g g' a^j$ and $\beta_j = \gamma_j * \beta_K * \beta'_K = \beta \beta' \chi^{-j}$.

□

In [11], the authors studied the representation theory of the Drinfel'd double of a family of Hopf algebras that generalize the Taft algebra. In their case, the order of the generating group-like element need not be the same as the order of the root of unity. They give a similar decomposition of tensor products as in Theorem IV.18. Although the algebras $H_{\mathcal{D}}$ generalize their Hopf algebras, Theorem IV.18 does not generalize their result since I require $|a| = |q|$. However, since G need not be cyclic, Theorem IV.18 generalizes Chen's result for Taft algebras.

CHAPTER V

CONCLUSION

In this dissertation I used Radford's method to construct simple modules for the Drinfel'd double of a graded Hopf algebra, to get information about the structure of these modules. I worked with two different classes of Hopf algebras: the restricted two-parameter quantum groups (of type A) defined by Benkart and Witherspoon in [6], and the rank one pointed Hopf algebras of nilpotent type introduced by Krop and Radford in [15].

For the two-parameter quantum groups, I presented necessary and sufficient conditions on the parameters r and s , for a simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ -module to be factored as the tensor product of a one-dimensional module with a module that is naturally a module for $\overline{\mathfrak{u}_{r,s}(\mathfrak{sl}_n)}$, the quotient of $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ by group-like central elements (Theorem II.13). In Chapter III, I introduced the code used in `SINGULAR::PLURAL` to construct simple $\mathfrak{u}_{r,s}(\mathfrak{sl}_3)$ -modules, and presented conjectures about bases and dimensions based on the computational results.

In Chapter IV, for $H_{\mathcal{D}}$ a rank one pointed Hopf algebra of nilpotent type, I gave necessary and sufficient conditions on \mathcal{D} for a simple $D(H_{\mathcal{D}})$ -module to factor as the tensor product of a one-dimensional module with a module that is naturally a module for $\overline{D(H_{\mathcal{D}})}$ (Theorem IV.10). Using this result, I studied the complete reducibility of the tensor product of two simple $D(H_{\mathcal{D}})$ -modules (Theorem IV.18). This result is a generalization of the work of Chen on the Drinfel'd double of the Taft algebra [7].

REFERENCES

- [1] N. Andruskiewitsch, H.-J. Schneider, On the classification of finite-dimensional pointed Hopf algebras, preprint (2005), arXiv.org:math/050215.
- [2] J. Apel, Gröbnerbasen in nichtkommutativen Algebren und Ihre Anwendung, Ph.D. thesis, Universität Leipzig (1988).
- [3] B. Bakalov, A. Kirillov, Jr., Lectures on tensor categories and modular functors, University Lecture Series, vol. 21, American Mathematical Society, Providence, RI (2001).
- [4] G. Benkart, S.-J. Kang, K.-H. Lee, On the centre of two-parameter quantum groups, Proc. Roy. Soc. Edinburgh Sect. A, 136 (3) (2006) 445–472.
- [5] G. Benkart, M. Pereira, S. Witherspoon, $\mathfrak{u}_q(\mathfrak{sl}_n)$ -modules constructed, in preparation.
- [6] G. Benkart, S. Witherspoon, Restricted two-parameter quantum groups, in: Representations of finite dimensional algebras and related topics in Lie theory and geometry, Fields Inst. Commun., vol. 40, Amer. Math. Soc., Providence, RI (2004), pp. 293–318.
- [7] H.-X. Chen, Irreducible representations of a class of quantum doubles, J. Algebra, 225 (1) (2000) 391–409.
- [8] V. Dobrev, Representations of quantum groups, in: Symmetries in Science, V (Lochau, 1990), Plenum, New York (1991), pp. 93–135.
- [9] V. Drinfel'd, Hopf algebras and the quantum Yang-Baxter equation, Dokl. Akad. Nauk SSSR, 283 (5) (1985) 1060–1064.

- [10] V. Drinfel'd, Quantum groups, in: Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI (1987), pp. 798–820.
- [11] K. Erdmann, E. Green, N. Snashall, R. Taillefer, Representation theory of the Drinfeld doubles of a family of Hopf algebras, *J. Pure Appl. Algebra*, 204 (2) (2006) 413–454.
- [12] G.-M. Greuel, V. Levandovskyy, H. Schönemann, SINGULAR::PLURAL 2.1, A computer algebra system for noncommutative polynomial algebras, Centre for Computer Algebra, University of Kaiserslautern (2003), <http://www.singular.uni-kl.de/plural>.
- [13] M. Jimbo, A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation, *Lett. Math. Phys.*, 10 (1) (1985) 63–69.
- [14] V. Kharchenko, A combinatorial approach to the quantification of Lie algebras, *Pacific J. Math.*, 203 (1) (2002) 191–233.
- [15] L. Krop, D. Radford, Finite-dimensional Hopf algebras of rank one in characteristic zero, *J. Algebra*, 302 (1) (2006) 214–230.
- [16] V. Levandovskyy, PBW bases, non-degeneracy conditions and applications, in: Representations of algebras and related topics, *Fields Inst. Commun.*, vol. 45, Amer. Math. Soc., Providence, RI (2005), pp. 229–246.
- [17] S. Majid, Doubles of quasitriangular Hopf algebras, *Comm. Algebra*, 19 (11) (1991) 3061–3073.
- [18] S. Montgomery, Hopf algebras and their actions on rings, *CBMS Regional Conference Series in Mathematics*, vol. 82, Published for the Conference Board of

the Mathematical Sciences, Washington, DC (1993).

- [19] D. Radford, On oriented quantum algebras derived from representations of the quantum double of a finite-dimensional Hopf algebra, *J. Algebra*, 270 (2) (2003) 670–695.
- [20] H.-J. Schneider, Normal basis and transitivity of crossed products for Hopf algebras, *J. Algebra*, 152 (2) (1992) 289–312.
- [21] E. Taft, The order of the antipode of finite-dimensional Hopf algebra, *Proc. Nat. Acad. Sci. U.S.A.*, 68 (1971) 2631–2633.
- [22] M. Takeuchi, A two-parameter quantization of $GL(n)$ (summary), *Proc. Japan Acad. Ser. A Math. Sci.*, 66 (5) (1990) 112–114.

VITA

Name: Mariana Pereira

Education:

M.S., Mathematics, University of Massachusetts, Amherst, Massachusetts, 2003

B.S., Mathematics, Universidad de la República, Montevideo, Uruguay, 2001

Ph.D., Mathematics, Texas A&M University, College Station, Texas, 2006

Address: c/o Dr. Sarah Witherspoon, Department of Mathematics, Texas A&M University, College Station, TX 77843-3368

e-mail: mpereira@math.tamu.edu